



**Europäisches
Patentamt**

**European
Patent Office**

**Office européen
des brevets**

REC'D 24 AUG 2004

WIPO

PCT

Bescheinigung

Certificate

Attestation

Die angehefteten Unterlagen stimmen mit der ursprünglich eingereichten Fassung der auf dem nächsten Blatt bezeichneten europäischen Patentanmeldung überein.

The attached documents are exact copies of the European patent application described on the following page, as originally filed.

Les documents fixés à cette attestation sont conformes à la version initialement déposée de la demande de brevet européen spécifiée à la page suivante.

Patentanmeldung Nr. Patent application No. Demande de brevet n°

03015400.9

**PRIORITY
DOCUMENT**
SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)

Der Präsident des Europäischen Patentamts;
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets
p.o.

R C van Dijk



Anmeldung Nr:
Application no.: 03015400.9
Demande no:

Anmeldetag:
Date of filing: 08.07.03
Date de dépôt:

Anmelder/Applicant(s)/Demandeur(s):

AXIOGENESIS AG
Josef-Stelzmann-Strasse 50,
Containergebäude
50931 Köln
ALLEMAGNE

Bezeichnung der Erfindung/Title of the invention/Titre de l'invention:
(Falls die Bezeichnung der Erfindung nicht angegeben ist, siehe Beschreibung.
If no title is shown please refer to the description.
Si aucun titre n'est indiqué se referer à la description.)

Secreted proteins as markers for cell differentiation

In Anspruch genommene Priorität(en) / Priority(ies) claimed /Priorité(s)
revendiquée(s)

Staat/Tag/Aktenzeichen/State/Date/File no./Pays/Date/Numéro de dépôt:

Internationale Patentklassifikation/International Patent Classification/
Classification internationale des brevets:

C12Q1/68

Am Anmeldetag benannte Vertragsstaaten/Contracting states designated at date of
filing/Etats contractants désignées lors du dépôt:

AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HU IE IT LU MC NL
PT RO SE SI SK TR LI

EPO - Munich
08. Juli 2003

Secreted Proteins As Markers For Cell Differentiation

5 Field of the invention

The present invention relates to the use of proteins capable of being secreted across the cell membrane as cell-type specific reporters during cell development. In detail, the present invention relates to embryonic stem (ES) cells stably transfected with a DNA construct comprising a DNA sequence encoding a secreted protein, which is operably linked to a cell-
10 and/or development-dependent promoter; a method for preparing such ES cells; a cell culture obtainable by culturing said ES cells; a method for the toxicological examination of substances using such cell cultures as well as methods for the identification of differentiation and growth factors for induction of selectively differentiated cells which can be used as a source for tissue grafts; a method for producing transgenic non-human mammals using said
15 ES cells; a transgenic non-human mammal obtainable by said method; and a method for examining stages of cellular development using cells of such a non-human mammal. The present invention also concerns assays to identify culture conditions and substances that promote the differentiation into a desired cell type, or to determine the toxicity of a compound for certain cell types. Those assays are particularly suited to give indications of possible
20 developmental and differentiation disturbances during embryogenesis.

Background art

Precursor cells have become a central interest in medical research. On the one hand precursors can replace cells that are senescent or damaged by injury or disease and on the other hand
25 these cells represent an ideal model for studying development and differentiation and the factors influencing these processes. Employing conventional cell lines for these studies has the disadvantage that individual cell lines may not be fully representative of the complex biology of an intact organism. Moreover, even repeating the tests in multiple cell lines does not reproduce or account for the complex interactions among cells and tissues that occurs in
30 an organism.

This is the main reason why toxicity testing in cell lines can not replace conventional animal tests. New chemical compounds are constantly being developed and tested on animals. In addition to industrial and household chemicals, a number of chemical compositions are developed each year for use as pharmaceuticals. Rules regarding the testing of potential
35 pharmaceuticals are promulgated by the Food and Drug Administration ("FDA"), which

currently requires comprehensive testing of toxicity, mutagenicity, and other effects in at least two species before a drug candidate can be entered into human clinical trials. Preclinical toxicity testing alone costs some hundreds of thousands of dollars. Despite this huge investment, almost one third of all prospective human therapeutics fail in the first phase of human clinical trials because of unexpected toxicity. It is clear that currently available toxicological screening assays do not detect all toxicities associated with human therapy or exposure to chemicals in the environment. Better means of screening potential therapeutics or chemicals in general for potential toxicity would reduce the cost and uncertainty of developing new therapeutics and materials, for example for use in medical devices or in other devices or goods humans are exposed to every day.

The detection of teratogenic and/or embryotoxic properties of chemical agents occurs presently by determination of the reproduction toxicity of test substances following single or multi-administrations to pregnant laboratory mammals and by tests of the embryotoxicity in the early stages of pregnancy. Furthermore, in vitro tests are performed with mammal embryos (Neubert and Merker, de Gruyter, Berlin-New York (1981)) and with embryonic organs for teratogenicity tests. These test procedures have however the disadvantage that they require the use of a large number of live mammals, in particular rats and mice. In vitro test procedures, in which primary cell cultures of limb buds (for example, "Limb Buds", Kochhar, Teratology 11 (1975), 273-287), or brain parts of embryonic rats (Flint and Orton, Toxicol. Appl. Pharmacol. 76 (1984), 383-395) or permanent cell lines of embryonic or adult mammal tissue, such as tumor cells of the ovary or embryonic palate cells are employed, do not fulfill, strictly speaking, the requirements which are imposed on the teratogenicity tests during the embryogenesis, namely giving indications of possible disgenesis or developmental disturbances.

Efforts have been made for a couple of years to employ permanent cultures of totipotent/pluripotent embryonic stem cells (ES cells) for the detection of embryotoxic and mutagenic substances; see, e.g. Laschinski et al., Reproductive Toxicol. 5 (1991), 57-65; Newall and Beedles, Toxicol. in Vitro 8 (1994), 697-701; Sehlmeier and Wobus, Mutation Res. 324 (1994), 69-76.

All those systems make use of the biological properties of the test ES cells or of a reporter system that has been introduced into the cells.

For example, US-A-6,498,018 describes a method for determining the effect of a biological agent by contacting a cell culture with a biological agent. The cell culture contains human multipotent CNS neural stem cells that are derived from primary CNS neural tissue and a culture medium with preselected growth factors. The read-out is provided by the effect of a biological agent on the presence or absence on a biological function or property ascribable to the cell culture. A major disadvantage of this system is the fact that particular biological functions or properties inherent to a certain culture of cells are difficult to measure and often involve the destruction of a large part of the culture in order to obtain enough material for the assay.

WO02/086073 discloses a method for the positive selection of neuronal cells differentiated from nuclear transfer embryonic stem cells by taking advantage of the neural stem cell marker nectin. This method is limited to neural cell types expressing nectin naturally.

US-A-6,007,993 describes an in vitro test procedure for the detection of chemically-induced embryotoxic (for example also teratogenic) effects based on differentiated pluripotent embryonic stem (ES) cells from the mouse and rat and using embryonic germ (EG) cells obtained established from primordial germ cells. Stable transgenic ES or EG stem cell clones are constructed, wherein a bacterial reporter gene, LacZ or the luciferase gene, is brought under the control of tissue-specific promoters or developmental control genes. Following differentiation of the ES cells in the presence of teratogenic substances into the different germination path derivatives, there occurs a differentiation-dependent expression in the cells, due to the activity of the tissue-specific promoters. The activation, repression or modulation of these tissue-specific genes is detected based on a reaction depending on the reporter gene employed, for example the X-Gal assay.

WO99/01552 discloses embryonic stem (ES) cells, which are transfected in a stable manner with a DNA construct encoding a non-cell damaging fluorescent protein and operatively linked thereto a cell- or development-dependent promoter. Also disclosed is a method for toxicological monitoring of substances using these ES cell cultures.

Although the above described methods employ semi-quantitative as well as relatively simple and robust assays, those assays usually involve the destruction of the tested cells. The preparation of a cell lysate required by many of the described conventional assay systems results in a high intra- as well as inter-assay variability. For repeated measurements during the course of differentiation many parallel cultures have to be assayed with the associated problems of variability. Laborious preparation of lysates also renders the assay less amenable

to high-throughput formats. Using fluorescent reporter proteins can overcome some of those problems. However, GFP expression has already shown some toxic effects (Liu et al. Biochem. Biophys. Res. Comm. 260 (1999), 712-717). Another disadvantage is the auto-fluorescence of aggregates of ES cells (embryoid bodies) which gives a high background in fluorescence measurements.

The technical problem of readily monitoring cells differentiating into a certain cell type and thereby providing a reliable and highly sensitive test system for toxicity screening, differentiation promoting agents and/or appropriate culture conditions, medium supplements, is achieved by providing the embodiments characterized in the claims, and described further below.

Summary of the invention

The present invention relates to a method of monitoring cell differentiation comprising culturing cells capable of differentiating into at least one particular cell type, wherein said cells contain at least one recombinant nucleic acid molecule comprising a reporter gene encoding a product that is secreted upon cell differentiation, under conditions allowing differentiation of the cells, or maintaining a non-human animal comprising such cells under appropriate conditions; and determining the amount or activity or amount of the reporter gene product either within a body fluid of said transgenic non-human animal or the cell culture medium or sample thereof.

The present invention also concerns the mentioned reporter gene constructs for monitoring cell differentiation, as well as cells, aggregates of at least one cell type, tissues comprising those cells or aggregates, organs comprising said cells or tissue, implants and transplants comprising said organ, all of which are characterized by comprising such a reporter gene construct.

Furthermore, the present invention relates to a non-human animal and a composition of matter both characterized by comprising a reporter gene construct, a cell, a cell aggregate, a tissue and/or an organ described herein.

In addition, the instant invention relates to an array or chip which contains the cells, cell aggregates or tissues of the invention attached to a solid support or suspended thereon as well as to an apparatus for analyzing said array or chip.

5 In particular, the present invention relates to test systems to identify substances that influence the differentiation of cells into certain cell types. Therefore, the present invention provides a method for obtaining and/or profiling a modulator of cell differentiation. This method comprises contacting a test sample containing a cell, cell aggregate, tissue, organ or a non-human transgenic animal as defined above with the substance to be tested; and then
10 determining the effect of the test substance on the amount of the reporter gene product or activity compared to a control sample or animal.

The test system provided by the present invention is useful for drug screening purposes. It is envisaged to obtain and manufacture drugs which promote or inhibit the formation of said
15 specific cell types using the method described above, wherein an enhanced or reduced level or activity of the reporter gene product is indicative for the drug. An enhanced or reduced level or activity of the reporter gene product, respectively, can also be indicative for an agent or compound which supports wound healing and/or healing of damaged tissue or toxicity of compounds, respectively.

20 The present invention additionally relates to a kit useful for conducting the methods described above, which contains, for example, a reporter gene construct and/or a cell as defined above, and standard compounds, like cell culture media, selection agents, detection agents for the reporter molecule and control samples.

25 Furthermore, methods of conducting a drug discovery and/or a target discovery business are disclosed. Both methods involve providing one or more assay systems described above for the identification of modulators of cell differentiation; and/or conducting therapeutic profiling of the modulators identified, or further analogs thereof, for efficacy and toxicity in the transgenic
30 animals provided by the present invention. The modulators identified using the methods provided by the invention are then either formulated in a pharmaceutical composition or licensed to a third party for further drug development and/or sales.

The present invention also relates to modulators of cell differentiation which have been identified using the methods of the invention as well as to pharmaceutical compositions comprising such modulators for use in the modulation of cell differentiation and the treatment of corresponding diseases.

5

Furthermore, the present invention relates to the use of a reporter gene construct, a cell, a cell aggregate, tissue, organ, implant or transplant, a non-human animal, a composition, an array or an apparatus provided herein in drug discovery, pharmacokinetic or pharmacological profiling.

10

Brief description of the figure

Figure 1: SEAP expression driven by the cardiac-specific α MHC promoter by cells differentiated from ES cells in the presence of retinoic acid.

15

ES cells transfected with the α -MHC-SEAP vector (containing the SEAP gene under the control of the heart-specific α MHC promoter) were differentiated in the presence of all-*trans* retinoic acid at concentrations indicated. After 13 days culture supernatants were assayed for SEAP activity. Shown are mean values of duplicate samples from which background SEAP activity was subtracted. Activities are expressed as percentage of SEAP activity in samples containing only solvent (DMSO) but no retinoic acid.

20

Definitions

For the purposes of this description, the term "stem cell" can refer to either stem cell or germ cell, for example embryonic stem (ES) and germ (EG), respectively. Minimally, a stem cell has the ability to proliferate and form cells of more than one different phenotype, and is also capable of self renewal-either as part of the same culture, or when cultured under different conditions. Embryonic stem cells are also typically telomerase positive and OCT-4 positive. Telomerase activity can be determined using TRAP activity assay (Kim et al., Science 266 (1997), 2011), using a commercially available kit (TRAPeze(R) XK Telomerase Detection Kit, Cat. s7707; Intergen Co., Purchase N.Y.; or TeloTAGGG(TM) Telomerase PCR ELISApplus, Cat. 2,013,89; Roche Diagnostics, Indianapolis). hTERT expression can also be evaluated at the mRNA level by RT-PCR. The LightCycler TeloTAGGG(TM) hTERT quantification kit (Cat. 3,012,344; Roche Diagnostics) is available commercially for research purposes.

30

In accordance with the present invention, the term embryonic stem (ES) cell includes any multi- or pluripotent stem cell derived from pre-embryonic, embryonic, or fetal tissue at any time after fertilization, and having the characteristic of being capable under appropriate conditions of producing progeny of several different cell types that are derivatives of all of the three germinal layers (endoderm, mesoderm, and ectoderm), according to a standard art-accepted test, such as the ability to form a teratoma in 8-12 week old SCID mice.

"Embryonic germ cells" or "EG cells" are cells derived from primordial germ cells. The term "embryonic germ cell" is used to describe cells of the present invention that exhibit an embryonic pluripotent cell phenotype. The terms "human embryonic germ cell (EG)" or "embryonic germ cell" can be used interchangeably herein to describe mammalian, preferably human cells, or cell lines thereof, of the present invention that exhibit a pluripotent embryonic stem cell phenotype as defined herein. Thus, EG cells are capable of differentiation into cells of ectodermal, endodermal, and mesodermal germ layers. EG cells can also be characterized by the presence or absence of markers associated with specific epitope sites identified by the binding of particular antibodies and the absence of certain markers as identified by the lack of binding of certain antibodies.

"Pluripotent" refers to cells that retain the developmental potential to differentiate into a wide range of cell lineages including the germ line. The terms "embryonic stem cell phenotype" and "embryonic stem-like cell" also are used interchangeably herein to describe cells that are undifferentiated and thus are pluripotent cells.

Included in the definition of ES cells are embryonic cells of various types, exemplified by human embryonic stem cells, described by Thomson et al. (Science 282 (1998), 1145); embryonic stem cells from other primates, such as Rhesus stem cells (Thomson et al., Proc. Natl. Acad. Sci. USA 92 (1995), 7844), marmoset stem cells (Thomson et al., Biol. Reprod. 55 (1996), 254) and human embryonic germ (hEG) cells (Shamblott et al., Proc. Natl. Acad. Sci. USA 95 (1998), 13726). Other types of pluripotent cells are also included in the term. Any cells of mammal origin that are capable of producing progeny that are derivatives of all three germinal layers are included, regardless of whether they were derived from embryonic tissue, fetal tissue, or other sources. The stem cells employed in accordance with the present invention that are preferably (but not always necessarily) karyotypically normal. However, it is preferred not to use ES cells that are derived from a malignant source.

"Feeder cells" or "feeders" are terms used to describe cells of one type that are co-cultured with cells of another type, to provide an environment in which the cells of the second type can grow. The feeder cells are optionally from a different species as the cells they are supporting.

5 For example, certain types of ES cells can be supported by primary mouse embryonic fibroblasts, immortalized mouse embryonic fibroblasts (such as murine STO cells, e.g., Martin and Evans, *Proc. Natl. Acad. Sci USA* 72 (1975), 1441-1445), or human fibroblast-like cells differentiated from human ES cells, as described later in this disclosure. The term "STO cell" refers to embryonic fibroblast mouse cells such as are commercially available and include
10 those deposited as ATCC CRL 1503.

The term "embryoid bodies" (EBs) is a term of art synonymous with "aggregate bodies". The terms refer to aggregates of differentiated and undifferentiated cells that appear when ES cells overgrow in monolayer cultures, or are maintained in suspension cultures. Embryoid bodies
15 are a mixture of different cell types, typically from several germ layers, distinguishable by morphological criteria; see also *infra*. As used herein, "embryoid body", "EB" or "EB cells" typically refers to a morphological structure comprised of a population of cells, the majority of which are derived from embryonic stem (ES) cells that have undergone differentiation. Under culture conditions suitable for EB formation (e.g., the removal of Leukemia inhibitory
20 factor or other, similar blocking factors), ES cells proliferate and form small mass of cells that begin to differentiate. In the first phase of differentiation, usually corresponding to about days 1-4 of differentiation for humans, the small mass of cells forms a layer of endodermal cells on the outer layer, and is considered a "simple embryoid body". In the second phase, usually corresponding to about days 3-20 post-differentiation for humans, "complex embryoid
25 bodies" are formed, which are characterized by extensive differentiation of ectodermal and mesodermal cells and derivative tissues. As used herein, the term "embryoid body" or "EB" encompasses both simple and complex embryoid bodies unless otherwise required by context. The determination of when embryoid bodies have formed in a culture of ES cells is routinely made by persons of skill in the art by, for example, visual inspection of the morphology.
30 Floating masses of about 20 cells or more are considered to be embryoid bodies; see, e.g., Schmitt et al., *Genes Dev.* 5 (1991), 728-740; Doetschman et al. *J. Embryol. Exp. Morph.* 87 (1985), 27-45. It is also understood that the term "embryoid body", "EB", or "EB cells" as used herein encompasses a population of cells, the majority of which being pluripotent cells capable of developing into different cellular lineages when cultured under appropriate

conditions. As used herein, the term also refers to equivalent structures derived from primordial germ cells, which are primitive cells extracted from embryonic gonadal regions; see, e.g., Shablott, et al. (1998), *supra*. Primordial germ cells, sometimes also referred to in the art as EG cells or embryonic germ cells, when treated with appropriate factors form pluripotent ES cells from which embryoid bodies can be derived; see, e.g., US Patent US-A-5,670,372; Shablott, et al., *supra*.

The terms "polynucleotide" and "nucleic acid molecule" refer to a polymer of nucleotides of any length. Included are genes and gene fragments, mRNA, tRNA, rRNA, ribozymes, cDNA, recombinant polynucleotides, branched polynucleotides, plasmids, vectors, isolated DNA and RNA, nucleic acid probes, and primers. As used in this disclosure, the term polynucleotides refer interchangeably to double- and single-stranded molecules. Unless otherwise specified or required, any embodiment of the invention that is a polynucleotide encompasses both a double-stranded form, and each of the two complementary single-stranded forms known or predicted to make up the double-stranded form. Included are nucleic acid analogs such as phosphoramidates and thiophosphoramidates.

A cell is said to be "genetically altered", "transfected", or "genetically transformed" when a polynucleotide has been transferred into the cell by any suitable means of artificial manipulation, or where the cell is a progeny of the originally altered cell that has inherited the polynucleotide. The polynucleotide will often comprise a transcribable sequence encoding a protein of interest, which enables the cell to express the protein at an elevated level. The genetic alteration is said to be "inheritable" if progeny of the altered cell have the same alteration.

A "regulatory sequence" or "control sequence" is a nucleotide sequence involved in an interaction of molecules that contributes to the functional regulation of a polynucleotide, such as replication, duplication, transcription, splicing, polyadenylation, translation, or degradation of the polynucleotide. Transcriptional control elements include promoters, enhancers, and repressors.

Particular gene sequences referred to as promoters, like the "αMHC" or "collagen" promoter, are polynucleotide sequences derived from the gene referred to that promote transcription of an operatively linked gene expression product. It is recognized that various portions of the

upstream and intron untranslated gene sequence may in some instances contribute to promoter activity, and that all or any subset of these portions may be present in the genetically engineered construct referred to. The promoter may be based on the gene sequence of any species having the gene, unless explicitly restricted, and may incorporate any additions, substitutions or deletions desirable, as long as the ability to promote transcription in the target tissue. Genetic constructs designed for treatment of humans typically comprise a segment that is at least 90% identical to a promoter sequence of a human gene. A particular sequence can be tested for activity and specificity, for example, by operatively linking to a reporter gene; see Figure 1.

According to the present invention, the term "cell- and/or development-dependent promoter" is intended to mean a promoter which displays its promoter activity only in particular cell types and/or only in particular stages of cellular development, both in cell cultures (embryoid bodies) and in transgenic non-human mammals derived from the ES cells according to the invention. In addition, any other known cell-specific promoter can be employed, e.g. for nerve cells, heart cells, neurons, glia cells, hematopoietic cells, endothelial cells, smooth muscle cells, skeletal muscle cells, cartilage cells, fibroblasts and epithelial cells.

Genetic elements are said to be "operatively linked" if they are in a structural relationship permitting them to operate in a manner according to their expected function. For instance, if a promoter helps initiate transcription of the coding sequence, the coding sequence can be referred to as operatively linked to (or under control of) the promoter. There may be intervening sequence between the promoter and coding region so long as this functional relationship is maintained.

In the context of encoding sequences, promoters, and other genetic elements, the term "heterologous" indicates that the element is derived from a genotypically distinct entity from that of the rest of the entity to which it is being compared. For example, a promoter or gene introduced by genetic engineering techniques into an animal of a different species is said to be a heterologous polynucleotide. An "endogenous" genetic element is an element that is in the same place in the chromosome where it occurs in nature, although other elements may be artificially introduced into a neighboring position.

The terms "polypeptide", "peptide" and "protein" are used interchangeably in this disclosure to refer to polymers of amino acids of any length. The polymer may comprise modified amino acids, it may be linear or branched, and it may be interrupted by non-amino acids.

- 5 If not stated otherwise the terms "compound", "substance" and "(chemical) composition" are used interchangeably herein and include but are not limited to therapeutic agents (or potential therapeutic agents), agents of known toxicities such as neurotoxins, hepatic toxins, toxins of hematopoietic cells, myotoxins, carcinogens, teratogens, or toxins to one or more reproductive organs. The chemical compositions can further be agricultural chemicals, such as
10 pesticides, fungicides, nematocides, and fertilizers, cosmetics, including so-called "cosmeceuticals", industrial wastes or by-products, or environmental contaminants. They can also be animal therapeutics or potential animal therapeutics.

Industrial products that can be tested with the methods of the present invention include bleaches, toilet, blocks, washing-up liquids, soap powders and liquids, fabric conditioners,
15 window, oven, floor, bathroom, kitchen and carpet cleaners, dishwater detergents and rinse aids, watersoftening agents, descalers, stain removers, polishes, oil products, paints, paint removers, glues, solvents, varnishes, air fresheners, moth balls and insecticides.

New ingredients for household products are constantly being developed and needed to be tested. For example, in recent years new enzymes (to digest stains) and "optical brighteners"
20 (which make washing appear whiter) have been developed for use in washing powders and liquids. New surfactants (which cut through grease to remove ingrained dirt) and chemical "builders" (which act as water softeners and enable surfactants to work more effectively) have been developed for use in washing powders and liquids, washing-up liquids and various cleaning agents. But also medical materials have to be tested, for example dental materials
25 such as new filling polymers, metal alloys, and bioactive ceramic. Furthermore, chemical compositions of any part of a device, such as catheters, electrodes, adhesives, paste, gel or cream may be tested with the method of the present invention in different concentrations and with different ingredients and impurities present.

- 30 As used herein, "molecular profile" or "profile" of a chemical composition or compound refers to a pattern of alterations in gene or protein expression, or both, in an ES cell, embryoid body, tissue, etc. contacted by the chemical composition compared to a like cell, embryoid body or tissue in contact only with culture medium.

Detailed description

Stem cells of various kinds have become an extremely attractive modality in regenerative medicine. They can be proliferated in culture, and then differentiated in vitro or in situ into the cell types needed. This plasticity makes them ideal models for toxicity testing as well as

5 investigation of pharmaceuticals such as tissue specific promoting factors. Particularly embryoid bodies which consist of different cell types of the three germ layers that interact with each other provide a highly sensitive test system. The ability of the stem cells to differentiate into the different cell types in the presence of a test substance may depend on the ability of the substance to promote differentiation and/or the toxicity of said substance.

10 This method of monitoring cell differentiation comprises:

- transfecting cells capable of differentiating into at least one particular cell type with at least one recombinant nucleic acid molecule comprising a reporter gene encoding a product that is secreted upon cell differentiation;
- introducing the cells into a non-human animal or culturing the cells under conditions
- 15 allowing differentiation of the cells; and
- determining the amount or activity of the reporter gene product either within a body fluid of said transgenic non-human animal or the cell culture medium.

The present invention is based on the use of a reporter gene that is secreted. Surprisingly it was found that secreted reporter molecules have certain advantages over the use of, for

20 example, a fluorescent reporter like Green Fluorescent Protein (GFP) and its derivatives. Liu et al. (Biochem. Biophys. Res. Comm. 260 (1999) 712-717) observed toxic effects when using a GFP-reporter. Such toxic effects can be avoided or at least minimized if a secreted reporter molecule is used, simply by exchanging the culturing medium on a regular basis and thereby preventing toxic concentration of the respective reporter molecule building up.

25 Also, a preferred method for the differentiation of cells is to grow them as embryoid bodies (EBs), which show a high autofluorescence rate even when the cells were lysed. This background can render the use of fluorescent reporter genes for the monitoring of cell differentiation quite inefficient.

Other routinely used reporter molecules like for example beta-galactosidase or luciferase

30 require the cells to be tested in a certain assay buffer which means that the cells have to be destroyed to determine the activity. The system according to the invention avoids this by employing a secreted reporter molecule so that assaying the culture medium of the differentiating cells or the body fluid an animal containing said cells is sufficient to monitor the reporter activity and thereby the differentiation.

For the transfection of the nucleic acid molecule encoding the reporter protein any standard method known to the person skilled in the art can be employed. Any suitable expression vector for this purpose can be used. Suitable viral vector systems for producing stem cells altered according to this invention can be prepared using commercially available virus components. The introduction of the vector construct or constructs into the cells occurs in a known manner, e.g. by transfection or transduction with the help of viral vectors. Transfection of vector plasmids into cells can be accomplished by, e.g., electroporation or lipofection, i.e. using lipid/DNA complexes. Exemplary is the formulation Lipofectamine 2000(TM), available from Gibco/Life Technologies. Another exemplary reagent is FuGENE(TM) 6 Transfection Reagent, a blend of lipids in non-liposomal form and other compounds in 80% ethanol, obtainable from Roche Diagnostics Corporation. . Viral vectors comprising effector genes are generally described in the publications referenced in the last section.

The cells to be transfected have to be able to differentiate into at least one particular cell type. This will be the case for the precursor cell for the desired cell type. Precursor cells include but are not limited to for example the human neuronal precursor cell line NTERA-2 cl.D1. (Leypoldt et al., J. Neurochem. 76 (2001), 806-814), the human retinal precursor cell line KGLDMSM (Ezeonu et al., DNA Cell Biol. 19 (2000), 527-537), a human leukemic cell line, PER-117, bearing the markers of a T-cell precursor phenotype (Kees, Blood 72 (1988), 1524-1529) or HL60, a promyelocytic leukemia cell line that differentiates to mature neutrophils (Fontana, Proc. Natl. Acad. Sci. USA 77 (1980), 3664-3668).

Since these precursor cell lines have only a limited range of developmental capacity it is preferred that multipotent or pluripotent cells are used. This is the case when said cells are or are derived from stem cells or primordial germ cells, which are preferred in the embodiments of the present invention.

The invention can be practiced using stem cells of any vertebrate species. Included are stem cells from humans; as well as non-human primates, domestic animals, livestock, and other non-human mammals. Preferred sources for stem cells are rodents, in particular mice. Amongst the stem cells suitable for use in this invention are primate pluripotent stem cells derived from tissue formed after gestation, such as a blastocyst, or fetal or embryonic tissue taken any time during gestation. Non-limiting examples are primary cultures or established lines of embryonic stem cells. The invention is also applicable to adult stem cells, for example multipotent adult progenitor cells (MAPCs); see, e.g., Reyes and Verfaillie, Ann. N Y Acad.

Sci. 938 (2001), 231-235 and Weissman and Anderson, *Annu. Rev. Cell. Dev. Biol.* 17 (2001), 387-403. It is also referred to the literature of Anderson et al., *Nat. Med.* 7 (2001), 393-395 and Prockop, *Science* 276 (1997), 71-74, wherein the extraction and culture of adult stem cells is described.

5

Media for isolating and propagating stem cells can have any of several different formulas, as long as the cells obtained have the desired characteristics, and can be propagated further. Suitable sources include Iscove's modified Dulbecco's medium (IMDM), Gibco, #12440-053; Dulbecco's modified Eagles medium (DMEM), Gibco #11965-092; Knockout
 10 Dulbecco's modified Eagles medium (KO DMEM), Gibco #10829-018; 200 mM L-glutamine, Gibco # 15039-027; non-essential amino acid solution, Gibco 11140-050; [beta]-mercaptoethanol, Sigma # M7522; human recombinant basic fibroblast growth factor (bFGF), Gibco # 13256-029. Exemplary serum-containing ES medium and conditions for culturing stem cells are known, and can be optimized appropriately according to the cell type. Media
 15 and culture techniques for particular cell types referred to in the previous section are provided in the references cited herein.

Because of their wide range of developmental capacity said stem cells are embryonic stem (ES) cells in a particularly preferred embodiment of this invention.

20 As mentioned before, several sources for ES cells are at the disposal of the skilled person of which human stem cells are preferred for most of the embodiments of the present invention. Human embryonic stem cells and their use for preparing different cell and tissue types are also described in *Reprod. Biomed. Online* 4 (2002), 58-63. Embryonic stem cells can be isolated from blastocysts of members of the primate species (Thomson et al., *Proc. Natl. Acad. Sci. USA* 92 (1995), 7844). Human Embryonic Germ (EG) cells can be prepared from
 25 primordial germ cells present in human fetal material taken about 8-11 weeks after the last menstrual period. Suitable preparation methods are described in Shamblott et al., *Proc. Natl. Acad. Sci. USA* 95 (1998), 13726. Methods for making cells that resemble embryonic stem cells or embryonic germ cells in morphology and pluripotency derived from primordial germ
 30 cells isolated from human embryonic tissue, such as from the gonadal ridges of human embryo, are described in US patent US-A-6,245,566.

Recently, it has been reported that exfoliated human deciduous tooth, a comparably very accessible tissue, contains multipotent stem cells that were identified to be a population of

highly proliferative, clonogenic cells capable of differentiating into a variety of cell types including neural cells, adipocytes, and odontoblasts; see Miura et al., *Proc. Natl. Acad. Sci. USA*, (2003), Apr 25; p S0027-8424. After in vivo transplantation, those cells were found to be able to induce bone formation, generate dentin, and survive in mouse brain along with expression of neural markers. Furthermore, multilineage potential of homozygous stem cells derived from metaphase II oocytes has been described in by Lin et al. in *Stem Cells* 21 (2003), 152-161. Various sources of precursor cells in postnatal muscles and the factors that may enhance stem cell participation in the formation of new skeletal and cardiac muscle in vivo are reviewed in Grounds et al. *J. Histochem. Cytochem.* 50 (2002), 589-610. Purification of rare Hematopoietic Stem Cell(s) (HSC) to homogeneity that home to bone marrow is described in US2003/0032185. These adult bone marrow cells are described to have tremendous differentiative capacity as they can also differentiate into epithelial cells of the liver, lung, gastro-intestinal tract, and skin. This finding may contribute to clinical treatment of genetic disease or tissue repair. Furthermore, techniques such as nuclear transfer for embryo reconstruction may be employed wherein diploid donor nuclei are transplanted into enucleated MII oocytes. This technology along with other procedures that aid in the establishment of customized embryonic stem (ES) cell lines that are genetically identical to those of the recipient have been reviewed by Colman and Kind, *Trends Biotechnol.* 18 (2000), 192-196.

20 The field of stem cell technology is being reviewed by Kiessling and Anderson, Harvard Medical School, in *Human Embryonic Stem Cells: An Introduction to the Science and Therapeutic Potential*; (2003) Jones and Bartlett Publishers; ISBN: 076372341X.

In order to avoid the use of for example human embryos as the donor for stem cells, which however seems to be justifiable at least under certain circumstances, it may even be possible to employ transgenic non-human animals, in particular mammals as source for embryonic stem cells. For example, compositions and methods for making transgenic swines to be used as xenograft donors is described in US patent US-A-5,523,226. Likewise, WO97/12035 describes methods of producing transgenic animals for xenotransplantation. Furthermore, immunologically compatible animal tissue, suitable for xenotransplantation into human patients, is described in WO01/88096. Methods for making embryonic germ cells from porcine are described for example in US patent US-A-6,545,199. Cells immunologically compatible with humans can also be employed for purposes of the present invention.

Stem cells can be propagated continuously in culture, using a combination of culture conditions that promote proliferation without promoting differentiation. Traditionally, stem cells are cultured on a layer of feeder cells, typically fibroblast type cells, often derived from embryonic or fetal tissue. The cell lines are plated to near confluence, usually irradiated to prevent proliferation, and then used to support when cultured in medium conditioned by certain cells (e.g. Koopman and Cotton, *Exp. Cell* 154 (1984), 233-242; Smith and Hooper, *Devel. Biol.* 121 (1987), 1-91), or by the exogenous addition of leukemia inhibitory factor (LIF). Such cells can be grown relatively indefinitely using the appropriate culture conditions without differentiation.

In the absence of feeder cells, exogenous leukemia inhibitory factor (LIF), or conditioned medium, ES or EG cells spontaneously differentiate into a wide variety of cell types, including cells found in each of the endoderm, mesoderm, and ectoderm germ layers. With the appropriate combinations of growth and differentiation factors, however, cell differentiation can be controlled. For example, mouse ES and EG cells can generate cells of the hematopoietic lineage in vitro (Keller et al., *Mol. Cell. Biol.* 13 (1993), 473-486; Palacios et al., *Proc. Natl. Acad. Sci. USA* 92 (1995), 7530-7534; Rich, *Blood* 86 (1995), 463-472). Additionally, mouse ES cells have been used to generate in vitro cultures of neurons (Bain et al., *Developmental Biology* 168 (1995), 342-357; Fraichard et al., *J. Cell Science* 108 (1995), 3161-3188), cardiomyocytes (heart muscle cells) (Klug et al., *Am. J. Physiol.* 269 (1995), H1913-H1921), skeletal muscle cells (Rohwedel et al., *Dev. Biol.* 164 (1994), 87-101), vascular cells (Wang et al., *Development* 114 (1992), 303-316), US patent US-A-5,773,255 relates to glucose-responsive insulin secreting pancreatic beta cell lines, US patent US-A-5,789,246 relates to hepatocyte precursor cells. Hepatic differentiation of murine embryonic stem cells is also described in Jones et al., *Exp. Cell Res.* 272 (2002), 15-22.

Other progenitors of interest include but are not limited to chondrocytes, osteoblasts, retinal pigment epithelial cells, fibroblasts, skin cells such as keratinocytes, dendritic cells, hair follicle cells, renal duct epithelial cells, smooth and skeletal muscle cells, testicular progenitors, and vascular endothelial cells.

In certain embodiments of the invention, differentiation is promoted by withdrawing one or more medium component(s) that promote(s) growth of undifferentiated cells, or act(s) as an inhibitor of differentiation. Examples of such components include certain growth factors, mitogens, leukocyte inhibitory factor (LIF), and basic fibroblast growth factor (bFGF).

Differentiation may also be promoted by adding a medium component that promotes differentiation towards the desired cell lineage, or inhibits the growth of cells with undesired characteristics.

- 5 In a preferred embodiment of the present invention, said reporter gene in the recombinant nucleic acid molecule described above is operably linked to at least one cell type specific regulatory sequence. This embodiment has the advantage of providing not only a monitoring system for differentiation per se but also allows observing the differentiation into a certain cell type. The regulatory sequence will drive the expression of the reporter molecule only in
10 those cells that acquire the desired cell type by differentiation. Usually, said regulatory sequence comprises such promoter and/or enhancer elements. In addition, or alternatively, a second reporter gene can be used which is under the control of said cell type specific regulatory sequence. The second reporter gene may be the same as used for the differentiation dependent expression or different, including conventional reporter genes such as green
15 fluorescent protein (GFP). Furthermore, the second reporter gene may be present in the mentioned recombinant nucleic acid molecule or on a different vector construct, the latter option requiring cotransfection of both molecules into the cells.

- In a further embodiment, the differentiated cells containing the recombinant nucleic acid molecule of the present invention, and optionally further foreign DNA such as vector
20 constructs described above, are enriched and separated from non-differentiated cells and/or from cells of non-desired cell types. Thus, in accordance with this invention, populations of differentiated cells are depleted of relatively undifferentiated cells and/or of cells of undesired cell types by using a selection system that is lethal to the undesired cells and cell types. This
25 can be achieved, for example, by expressing a selectable marker gene under the control of a regulatory sequence that causes the gene to be preferentially expressed in the desired cell type and/or at a certain stage of development, which renders cells of a specific cell type resistant to a lethal effect of an external agent. To accomplish this, the cells are genetically altered with a recombinant nucleic acid molecule comprising the selectable marker gene before the process
30 used to differentiate the cells into the desired lineage. The selection system may be present in the recombinant nucleic acid molecule of the present invention or on a different vector construct which is used for cotransfection.

The cell types, which the differentiation can be directed to, include connecting fibroblasts, stromal cells, endothelial cells, glia cells, neural cells, neuronal cells, hematopoietic cells, smooth muscle cells, skeletal muscle cells, epithelial cells, and cardiac cells. To monitor differentiation into these particular cell types, regulatory sequences have to be operatively
 5 linked to the reporter gene sequence that confer an expression of the reporter which is specific for said cell types.

In a preferred embodiment of the method according to the invention said promoter or enhancer is selected from the group consisting of α MHC, MLC2V, VE-cadherin, Tie-2, Flk-1, Flt-1, GFAP, alpha-1-tubulin and collagen 2 promoter or enhancer. Whereas the pro-alpha 1
 10 (II) collagen chain (collagen 2) promoter/enhancer (Zhou et al., J. Cell Sci. 108 (1995), 3677-3684) is specific for chondrocytes, the expression in endothelial cells can be driven by the VE-cad promoter (Gory et al., Blood 93 (1999), 184-192), the Flt-1 promoter (Quinn et al., Biochem. Biophys. Res. Commun. 276 (2000), 1089-1099), the Tie-2 (Schlaeger et al., Proc.
 15 Natl. Acad. Sci. USA 94 (1997), 3058-3063) and the Flk-1 promoter/enhancer (Kappel et al., Blood 93 (1999), 4284-4292), respectively. The alpha-1-tubulin promoter (Gloster et al., J. Neurosci. 14 (1994), 7319-7330) and the glial fibrillary acidic protein (GFAP) promoter (Besnard et al., J. Biol. Chem. 266 (1991), 18877-18883) are specific for neural cells. The smooth muscle heavy chain (SMHC) minimal promoter is specific for smooth muscles
 20 (Kallmeier et al., J. Biol. Chem. 270(1995), 30949-30957). An HSV-1 vector containing the rat tyrosine hydroxylase promoter for use in both long-term and cell type-specific expression in the midbrain is described by Song et al., in J. Neurochem. 68 (1997), 792-803. Cardiac specific promoters such as alpha-myosin heavy chain promoter (α MHC) and the myosin light chain-2v (MLC2v) promoter, the latter being specific for ventricular heart muscles are
 25 described further below.

Further examples for tissue specific promoters are those, which are active in hematopoietic cells, cartilage cells or epidermal cells as well as insulin secreting β -cells.

"Tissue specific" is to be subsumed under the term "cell specific". Regulatory sequences of the above described cell types and other cell type specific promoters can be obtained from the
 30 literature; see, e.g., "medline" and NCBI.

Cardiovascular diseases are still among the most frequent causes of death in the Western industrial countries. Only through intensive basic research in this field, the pathophysiological causes can be found, and new therapeutic approaches can be devised, and toxicological

changes described. For studying the pathogenesis of cardiovascular diseases and for testing new pharmacological and toxicological substances, models are needed which, on one hand, can be transferred to humans and, on the other hand, can replace the tedious and cost-intensive animal models. Still in the year 1991, over 2 million animals have been employed for animal experiments in the federal states of the former Western Germany alone.

One starting point of pharmacological and toxicological research which has recently grown more and more important is heart differentiation. From the stereotypically proceeding heart cell development, conclusions can be drawn to pathological and toxicological changes of cardiomyocytes. Thus, for example, it is known that the receptor state and the intracellular signal cascades are disturbed in cardiac hypertrophy (Yamazaki et al., *J. Mol. Cell Cardiol.* 27 (1995), 133-140) and heart insufficiency (Johnson et al., *Biochem. Pharmacol.* 45 (1993), 2365-2372). These pathologically altered cardiomyocytes are in part again similar to heart cells of early differentiation stages. Thus, besides toxicological tests assays for the identification and isolation of substances that are capable of promoting cardiac remodeling are needed, which can be used for therapeutic interventions in the treatment of patients with cardiac dysfunction characterized by, for example, pathological hypertrophy and fibrosis.

However, the examinations necessary for elucidating the properties of heart cells in early differentiation stages are technically difficult to perform with live animals and, if they can be performed at all, it is only in very complicated studies: On day 12-13, at the earliest, it is possible to prepare cardiomyocytes from a murine embryo, but such cells no longer correspond to an early cardiac differentiation stage. A detailed analysis of receptor expression during various differentiation stages requires a very high expense of animal material and is technically difficult to perform as set forth above. Similarly, the observation of the development of a relatively undifferentiated heart cell over several days or weeks is not possible with an animal model. In order to test the use of novel therapeutical agents, e.g., inotropic substances or antiarrhythmics or toxic substances, e.g., heavy metals or retinoids, an invasive monitoring of animals, e.g., swine, must be performed for weeks in animal experiments.

Now, it has been the object of the present invention to provide a cell based assay which enables a simple characterization of the effects compounds on cardiomyogenesis and allows for functional examinations rather than being based on reporter genes such as Lac-Z (Niwa et al., *Gene* 108 (1991), 193-199; Wobus et al., *J. Mol. Cell Cardiol.* 29 (1997), 1525-1539;

Metzger et al., *Circ. Res.* 78 (1996), 547-552), as with prior methods, the expression of which genes can be detected only after cell fixation and by means of a specific substrate.

Surprisingly, it has been found that ES cells can be stably transfected with a DNA construct in which a gene encoding a secreted reporter protein is coupled with a cardiomyocyte specific cell- and development-dependent promoter; see the examples and the vector sequence depicted in SEQ ID NO: 3. This construct is integrated in the native DNA and after the specific activation of intracellular signals, the promoter is activated and the reporter protein expressed and secreted into the medium. Thus, ES cells which activate a cell-specific transcription factor at a certain point of differentiation could be recognized by determining the level of reporter protein or its activity in the cell culture medium.

Thus, in a particular preferred embodiment of the invention, said differentiating cell type are cardiomyocytes. For this embodiment, said cell type specific regulatory sequence is preferably atrial and/or ventricular specific. Corresponding regulatory sequences, i.e. cardiac specific promoters are described in the prior art; see also supra. For example Nkx-2.5 specific for very early cardiomyocytes and mesodermal precursor cells, respectively, (Lints et al., *Development* 119 (1993), 419-431); human-cardiac- α -actin specific for heart tissue, (Sartorelli et al., *Genes Dev.* 4 (1990), 1811-1822), and MLC-2V specific for ventricular heart muscle cells (O'Brien et al., *Proc. Natl. Acad. Sci. USA.* 90 (1993), 5157-5161; Lee et al., *Mol. Cell Biol.* 14 (1994), 1220-1229; Franz et al., *Circ Res.* 73 (1993), 629-638 and WO-A-96/16163). The cardiac specific alpha-myosin heavy chain promoter is described in Palermo et al., *Cell. Mol. Biol. Res.* 41 (1995), 501-519 and Gulick et al., *J. Biol. Chem.* 266 (1991), 9180-91855. The expression of the atrial specific myosin heavy chain AMHC1 and the establishment of anteroposterior polarity in the developing chicken heart is described in Yutzey et al., *Development* 120 (1994), 871-883.

Since the detection is aided by using a reporter molecule for which a robust, sensitive and specific assay exists, a particularly preferred embodiment is the method according to the invention, wherein said reporter gene product is secreted alkaline phosphatase (SEAP), alpha-amylase or invertase.

SEAP is heat-resistant and not inhibited by L-homoarginine. Further advantages of this reporter molecule are its low toxicity and the fact that the protein levels that are secreted correspond directly to the amount of intracellular mRNA. Thus the SEAP reporter assay

provides a direct measurement for the cell type-specific expression and thereby the differentiation. It is also extremely sensitive: it is possible to detect 10^{-13} g SEAP protein (Yang et al. *Biotechniques* 23 (1997), 1110-1114). This makes it possible to detect differentiation into cell types that are rare.

- 5 The use of alpha-amylase as reporter molecule is described in for example WO98/49320. This reporter molecule is particularly attractive because the enzyme is extremely stable over a wide range of conditions. The measurement of alpha-amylase activity is simple, quantitative, sensitive, safe and inexpensive. It was also shown to be compatible with a SEAP reporter system (Schlatter et al., *Gene* 282 (2002), 19-31). The use of invertase as a marker protein is
10 described for example in Mace et al., *Virology* 188 (1992), 869-874.

- Reporter gene constructs comprising inter alia secreted alkaline phosphatase as the reporter molecule and, which therefore can be used as a source for preparing vector constructs according to the present invention are described for example in WO00/34435 and Wang et al.,
15 *Gene* 279 (2001), 99-108, the latter describing murine SEAP. Detection of SEAP level can be performed as described in the Examples and in the prior art; see, e.g., Yang et al. (1997) and Wang et al. (2001), *supra*. The use of human secreted alkaline phosphatase in an in vivo mouse reporter gene model to monitor ovarian tumor growth and response to therapeutics has been described in Nilsson et al., *Cancer Chemother. Pharmacol.* 49 (2002), 93-100. This
20 model can be adapted to the embodiments of the present invention. Engineered secreted alkaline phosphatase reporter genes being capable of being expressed in a mammal for extended periods of time and detection of the alkaline phosphatase activity by measuring the protein levels or AP activity of a serum or tissue sample from a mammal containing the cell with the transgene, or measuring the media from a cultured cell containing the transgene are
25 also described in WO02/095068.

Reporter activity or protein amount, in particular SEAP activity, can be detected in any biological fluid such as breast milk, amniotic fluid, seminal plasma, serum and urine.

- Since not all molecules that can be used as a reporter are secreted naturally, the nucleic acid
30 molecule introduced into the cells can be genetically engineered to encode a protein that is excreted. Particularly preferred is the method described above, wherein said reporter gene product comprises a secretory leader sequence. Examples for secretory leader sequences are but are not limited to the adenovirus tripartite leader sequence (Chu et al., *Biotechniques* 5 (1995), 890-896), the immunoglobulin secretion signal (Schlatter et al., *Gene* 282 (2002), 19-

31), the SUC2 signal sequence as well as the human interleukin 2 receptor alpha chain signal sequence (Kamiya et al., Tohoku J. Exp. Med. 180 (1996), 297-308).

5 In order to select those cells which contain the recombinant nucleic acid molecule according to the invention, and optionally other vector constructs such as those described hereinbefore, said recombinant nucleic acid molecule and vector constructs, respectively, preferably further comprise a selectable marker expressed by multi- or pluripotent cells. Selectable markers are known to persons skilled in the art and include for example nucleoside and aminoglycoside-antibiotic-resistance genes conferring resistance to, e.g. puromycin, neomycin, or
10 hygromycin. Further examples for resistance genes are dehydrofolate-reductase, which confers a resistance against aminopterin and methotrexate, as well as multi drug resistance genes, which confer a resistance against a number of antibiotics, e.g. against vinblastin, doxorubicin and actinomycin D; see, e.g., Blau et al., Blood 89 (1997), 146-154, which describes the use of the multiple drug resistance gene 1 (MDR1) gene as a dominant
15 selectable marker.

In a particularly preferred embodiment of the present invention, said selectable marker confers resistance to puromycin. Puromycin is particularly suited for the fast elimination of non-cardiac cells in adherent culture of transgenic EBs. Furthermore, drug selection of cardiac
20 cells can be implemented entirely in the suspension culture of transgenic EBs. Hence, it could also be shown that purified ES derived cardiomyocytes survive much longer in culture than untreated counterparts. Moreover, the elimination of undifferentiated ES cells during drug selection process has itself been shown to have clear positive effect on viability and longevity of such differentiated ES derived cells as cardiomyocytes. In addition, it could be surprisingly
25 shown that the release from surrounding non-differentiated cells induces proliferation of cardiomyocytes. Thus, the drug selection possesses both purifying and multiplying effect; see, e.g., WO02/051987. Thus, the selection does not only provide the means for eliminating cells which can not be monitored since they do not carry the reporter gene, is also allows the elimination of cells that did not differentiate into the desired cell type provided that the
30 reporter gene and the selectable marker gene are driven by the same, cell type-specific regulatory sequence.

The method according to the invention, wherein said cells form cell aggregates or tissue-like aggregates derived from different ES cell types is a preferred embodiment and particularly

preferred are said cells forming embryoid bodies (EBs). As mentioned before, embryoid bodies represent a complex group of cells differentiating into different tissues. In one embodiment, the cells within an embryoid body are substantially synchronized for their differentiation. Accordingly, at known intervals, the majority of the synchronized cells differentiate into the three embryonic germ layers and further differentiate into multiple tissue types, such as cartilage, bone, smooth and striated muscle, and neural tissue, including embryonic ganglia; see also Snodgrass et al., "Embryonic Stem Cells: Research and Clinical Potentials" in Smith and Sacher, eds. *Peripheral Blood Stem Cells* American Association of Blood Banks, Bethesda MD (1993). Thus, the cells within embryoid bodies provide a much closer model to the complexity of whole organisms than do traditional single cell or yeast assays, while still avoiding the cost and difficulties associated with the use of mice and larger mammals. Moreover, the recent availability of human embryoid bodies improves the predictive abilities of the invention by providing an even closer vehicle for modeling toxicity and identification of drugs useful for the treatment of heart disorders in human organ systems, and in humans.

The embryoid body of the invention comprises a cell population, the majority of which being pluripotent cells capable of developing into different cellular lineages when cultured under appropriate conditions. It is preferred that the embryoid body comprises at least 51% pluripotent cells derived from totipotent ES cells. More preferably, the embryoid body comprises at least 75% pluripotent cells derived from totipotent ES cells. And still more preferably, the embryoid body comprises at least 95% pluripotent cells derived from totipotent ES cells.

In its simplest form, the method of creating a molecular profile according to the present invention involves contacting embryoid bodies with a chemical composition of interest, and then determining the alterations in reporter activity or amount of reporter protein, or both, in the embryoid body exposed to the chemical composition (the "test embryoid body") compared to a embryoid body which was not exposed to the agent (a "control embryoid body").

Thus, the method of the present invention can be performed such that it allows self-assembly of the cells into the aggregates or tissue-like structures. The term "embryoid bodies" (EBs) is a term of art synonymous with "aggregate bodies". The terms refer to aggregates of differentiated and undifferentiated cells that appear when ES cells overgrow in monolayer cultures, or are maintained in suspension cultures. Embryoid bodies are a mixture of different

cell types, typically from several germ layers, distinguishable by morphological criteria. WO02/051987 describes a protocol to obtain embryoid bodies. The manufacturing takes place preferably with the "hanging drop" method or by methylcellulose culture (Wobus et al., Differentiation 48 (1991), 172-182).

5 Alternatively to this, spinner flasks (stirring cultures) can be used as culture method. Therefore, the undifferentiated ES-cells are introduced into stirring cultures and are mixed permanently according to an established procedure. Therefore, 10 million ES-cells are introduced into 150 ml medium with 20 % FCS and are stirred constantly with the rate of 20 rpm., wherein the direction of the stirring motion is changed regularly. 24 hours after
 10 introduction of the ES-cells an extra 100 ml medium with serum is added and thereupon 100 - 150 ml of the medium is exchanged every day (Wartenberg et al., FASEB J. 15 (2001), 995-1005). Under these culture conditions large amounts of ES-cell-derived cells, i.e. cardiomyocytes, endothelial cells, neurons etc. depending on the composition of the medium can be obtained. The cells are selected by means of the resistance gene either still within the
 15 stirring culture or after plating, respectively.

In a particular preferred embodiment of the present invention, embryoid bodies are prepared according a recent developed "mass culture" system employed in the appended examples and described in detail in applicant's co-pending application "Novel method for the preparation of Embryoid Bodies (EBs) and uses thereof", Attorneys docket no: AX02A15/P-EP, filed on the
 20 same day as the present application, i.e. on July 8, 2003.

The embryoid bodies used to test the chemical composition can be of any vertebrate species. The choice of the particular species from which the embryoid body is derived will typically reflect a balance of several factors. First, depending on the purpose of the study, one or more
 25 species may be of particular interest. For example, human embryoid bodies will be of particular interest for use with compositions being tested as potential human therapeutics but also for toxicological tests for substances including industrial chemicals, while equine, feline, bovine, porcine, caprine, canine, or sheep embryoid bodies may be of more interest for a potential veterinary therapeutic. Embryoid bodies of other species commonly used in
 30 preclinical testing, such as guinea pigs, mice, rat, rabbits, pigs, and dogs, are also preferred. Typically, embryoid bodies of these species will be used for "first pass" screening, or where detailed information on toxicity in humans is not needed, or where a result in a murine or other one of these laboratory species has been correlated to a known toxicity or other effect in humans. Furthermore, with respect to human therapeutics, regulatory agencies generally

require animal data before human trials can begin; it will generally be desirable to use embryoid bodies of species which will be used in the preclinical animal studies. The results of toxicity testing in the embryoid bodies can then guide the researcher on the degree and type of toxicity to anticipate during the animal trials. Certain animal species are known in the art to be better models of human toxicity of different types than are others, and species also differ in their ability to metabolize drugs; see, e.g., Williams, *Environ. Health Perspect.* 22 (1978), 133-138; Duncan, *Adv. Sci.* 23 (1967), 537-541. Thus, the particular species preferred for use in a particular preclinical toxicity study may vary according to the intended use of the drug candidate. For example, a species which provide a suitable model for a drug intended to affect the reproductive system may not be as suitable a model for a drug intended to affect the nervous system. Criteria for selecting appropriate species for preclinical testing are well known in the art.

Once an embryoid body culture has been initiated, it can be contacted with a chemical composition. Conveniently, the chemical composition is in an aqueous solution, preferably in a solvent conventionally used in cell culture, for example DMSO, and is introduced to the culture medium; see also the examples. The introduction can be by any convenient means, but will usually be by means of a pipette, a micropipettor, or a syringe. In some applications, such as high throughput screening, the chemical compositions will be introduced by automated means, such as automated pipetting systems, which may be on robotic arms. Chemical compositions can also be introduced into the medium as in powder or solid forms, with or without pharmaceutical excipients, binders, and other materials commonly used in pharmaceutical compositions, or with other carriers which might be employed in the intended use. For example, chemical compositions intended for use as agricultural chemicals or as petrochemical agents can be introduced into the medium by themselves to test the toxicity of those chemicals or agents, or introduced in combination with other materials with which they might be used or which might be found in the environment, to determine if the combination of the chemicals or agents has a synergistic effect. Typically, the cultures will be shaken at least briefly after introduction of a chemical composition to ensure the composition is dispersed throughout the medium.

The time as which a chemical composition is added to the culture is within the discretion of the practitioner and will vary with the particular study objective. Conveniently, the chemical composition will be added as soon as the embryoid body develops from the stem cells,

permitting the determination of the alteration in protein or gene expression on the development of all the tissues of the embryoid body. It may be of interest, however, to focus the study on the effect of the composition on a particular tissue type. As previously noted, individual tissues, such as muscle, nervous, and hepatic tissue, are known to develop at specific times after the embryoid body has formed. Addition of the chemical composition can therefore be staged to occur at the time the tissue of interest commences developing, or at a chosen time after commencement of that development, in order to observe the effect on altering gene or protein expression in the tissue of interest.

- 10 Different amounts of a chemical composition will be used to contact an embryoid body depending on the amount of information known about the toxicity of that composition, the purposes of the study, the time available, and the resources of the practitioner. A chemical composition can be administered at just one concentration, particularly where other studies or past work or field experience with the compound have indicated that a particular concentration is the one which is most commonly found in the body. More commonly, the chemical composition will be added in different concentrations to cultures of embryoid bodies run in parallel, so that the effects of the concentration differences on gene or protein expression and, hence, the differences in toxicity of the composition at different concentrations, can be assessed. Typically, for example, the chemical composition will be added at a normal or medium concentration, and bracketed by twofold or fivefold increases and decreases in concentration, depending on the degree of precision desired.

- Where the composition is one of unknown toxicity, a preliminary study is conveniently first performed to determine the concentration ranges at which the composition will be tested. A variety of procedures for determining concentration dosages are known in the art. One common procedure, for example, is to determine the dosage at which the agent is directly toxic. The practitioner then reduces the dose by one half and performs a dosing study, typically by administering the agent of interest at fivefold or twofold dilutions of concentration to parallel cultures of cells of the type of interest. For environmental contaminants, the composition will usually also be tested at the concentration at which it is found in the environment. For agricultural chemicals, such as pesticides which leave residues on foodstuffs, the agent will usually be tested at the concentration at which the residue is found, although it will likely be tested at other concentrations as well. Thus, the dilution of test compounds can be done by making in separated tubes a series of dilution of 50 or 100

fold concentrated compounds in DMSO. One or two μ l of each dilution are distributed in each well before cell suspension distribution. Particularly preferred are stock and dilution series as described for retinoic acid in example 3.

- 5 The above considerations with respect to contacting the compounds with the EBs, contacting time, etc, also apply to the assays of the invention performed on e.g. ES cells, tissue and non-human animals, if applicable.

10 The present invention also relates to reporter gene constructs for monitoring cell differentiation comprising the recombinant nucleic acid molecule defined previously as well as to cells capable of differentiating into at least one particular cell type, that contain said reporter gene construct, or are obtainable by the methods described herein. Likewise, cell aggregates, tissues, organs, implants or transplants comprising said reporter gene construct, obtainable by the methods described herein or constituted from said cells, cell aggregates,
15 tissues, organs, implants or transplants are subject of the invention.

In a still further embodiment, the present invention relates to a transgenic non-human animal which can be generated from the mentioned ES cells and ES cell derived cell types and cell aggregates; see supra. A method for the production of a transgenic non-human animal, which
20 is encompassed by the present invention, for example transgenic mouse, comprises introduction of a reporter gene as described above into a germ cell, an embryonic cell, stem cell or an egg or a cell derived therefrom. The non-human animal can be used in accordance with a screening method of the invention described herein. Production of transgenic embryos and screening of those can be performed, e.g., as described by A. L. Joyner Ed., Gene
25 Targeting, A Practical Approach (1993), Oxford University Press. A general method for making transgenic non-human animals is described in the art, see for example WO 94/24274. For making transgenic non-human organisms (which include homologously targeted non-human animals), embryonal stem cells (ES cells) are preferred. Murine ES cells, such as AB-1 line grown on mitotically inactive SNL76/7 cell feeder layers (McMahon and Bradley, Cell
30 62 (1990), 1073-1085) essentially as described (Robertson, in Teratocarcinomas and Embryonic Stem Cells: A Practical Approach. E. J. Robertson, ed. (Oxford: IRL Press) (1987), 71-112) may be used for homologous gene targeting. Other suitable ES lines include, but are not limited to, the E14 line (Hooper et al., Nature 326 (1987), 292-295), the D3 line (Doetschman et al., J. Embryol. Exp. Morph. 87 (1985), 27-45), the CCE line (Robertson et

al., Nature 323 (1986), 445-448), the AK-7 line (Zhuang et al., Cell 77 (1994), 875-884) and others; see also supra. The success of generating a mouse line from ES cells bearing a specific targeted mutation depends on the pluripotency of the ES cells (i.e., their ability, once injected into a host developing embryo, such as a blastocyst or morula, to participate in embryogenesis and contribute to the germ cells of the resulting animal). The blastocysts containing the injected ES cells are allowed to develop in the uteri of pseudopregnant nonhuman females and are born as chimeric mice. The resultant transgenic mice are backcrossed and screened for the presence of the correctly targeted transgene (s) by PCR or Southern blot analysis on tail biopsy DNA of offspring so as to identify transgenic mice for the reporter locus/loci. For example, transient expression of a recombinase gene directly introduced into fertilized eggs by pronuclear injection provides a fast and efficient method for generating mutant mice with desired deletions or translocations in target genes. Efficient removal of loxP-flanked DNA sequences in a gene-targeted locus by transient expression of Cre recombinase in fertilized eggs is described by Sunaga et al., Mol. Reprod. Dev. 46 (1997), 109-113.

Methods for producing transgenic flies, such as *Drosophila melanogaster* are also described in the art, see for example US-A-4,670,388, Brand and Perrimon, Development 118 (1993), 401-415; and Phelps and Brand, Methods 14 (1998), 367-379. Transgenic worms such as *C. elegans* can be generated as described in Mello et al., EMBO J. 10 (1991), 3959-3970, Plasterk, Methods Cell. Biol. 48 (1995), 59-80.

All of the embodiments above can not only be used to monitor cell differentiation and optimize culturing conditions using modulators in their pure forms but the effective agent(s) might also be contained in a composition. Hence, compositions of matter comprising any one of those recombinant nucleic acid molecules, cells, cell aggregates, or tissues of the present invention as described herein are encompassed in the scope of the present invention. As described herein, those compositions and methods of the invention can be used for a variety of purposes, for example for analyzing early steps of tissue formation during embryonic development or the influence of factors and compounds on this process.

In a particularly preferred embodiment, the present invention relates to arrays and chips comprising a solid support and attached thereto or suspended thereon cells, cell aggregates or tissue obtained by the method of the present invention or being in the differentiation process. Such arrays generally consist of a substrate of glass, plastic or silicon on which the test cells

aggregates or tissues are deposited in a particular pattern. Depending on the type of array, these might additionally be covered by a conductor, e.g. gold, platinum, indium-tin-oxide, iridium, etc., which allows a direct measurement by employing the conductivity of cells. The use of such planar microelectrode arrays for cultured cells and cell aggregates as biosensors is of particular interest.

Furthermore the invention relates to an apparatus for analyzing said array, which will be able to measure the particular read-out of the assay employed, e.g. SEAP reaction product, in a quantitative manner and assign the results to the position on the array.

The system for monitoring cell differentiation according to the present invention can also be employed to identify and characterize substances that modify the differentiation process. Populations of differentiated cells and tissue suitable for human administration have to be identified after the differentiation process to avoid administering undifferentiated cells with high tumorigenic potential or cells differentiated into cells that are not needed for the particular therapeutic purpose. The positive identification of cells differentiated into a certain cell type allows also the optimization of culturing conditions that is necessary to obtain the relatively high numbers of differentiated cells needed for therapeutic purposes. Thus, there is a need to identify conditions for culturing ES cells that drive their differentiation into the desired direction.

Accordingly, a preferred embodiment of the invention is a method of obtaining and/or profiling a modulator of cell differentiation comprising:

- contacting a test sample comprising a cell, cell aggregate, tissue or organ or a non-human animal as defined above with a test substance; and
- determining the effect of the test substance on the amount of the reporter gene product or activity compared to a control sample or animal.

In particular, the methods of the invention can be used for toxicological, embryotoxic, mutagenic, and/or teratogenic in vitro tests; see supra. Another important aspect of the present invention is therefore a method of determining toxicity, preferably teratogenicity, embryotoxicity, chronic or acute toxicity of a compound comprising the steps of the methods described herein, wherein a reduced or enhanced level or activity of said reporter gene product is indicative for the toxicity of the compound.

The assays may be simple "yes/no" assays to determine whether there is a change in expression of the secreted reporter protein by detecting reporter, e.g., enzyme activity or the amount of reporter protein. The assay may be made quantitative by comparing the amount of activity and/or amount of reporter in a test sample with the levels of activity and/or amount in a standard sample. As described in the examples, the test compound or a plurality of test compounds are subjected to the test cell, preferably embryoid body in different concentrations or dilution series, preferably at doses that correspond to physiological levels of the corresponding type of test compounds; see also example 3. It is thus also possible to easily generate compound profiles in purpose similar to those described in WO00/34525. For example, two or more assays may be used, wherein for each assay the test cells or, e.g., embryoid bodies comprise recombinant DNA molecules with different cell- and/or development-specific promoters operably linked to the same or different reporters, preferably wherein each reporter is a secreted protein, e.g., enzyme, most preferably wherein one or each reporter is SEAP. Those assays can be performed in parallel or subsequently; or the results of one assay may be compared with results of a corresponding assay performed elsewhere. Likewise, it is envisaged to provide and use cells which comprise two of the mentioned recombinant molecules but with different promoters and reporters as mentioned above. Thus, a molecular profile of a test chemical composition can be established by detecting the alterations in reporter gene activity or amount of reporter protein in, for example, embryoid bodies contacted by the test chemical composition as described in previous sections. Once the molecular profile of the test composition is determined, it can be compared to that of a chemical composition with predetermined toxicities or, preferably, to a library of molecular profiles of chemical compositions with predetermined toxicities. The outcome of such comparison provides information for one to predict the likelihood of whether the test composition is toxic, what type of toxicities, and how toxic it would be as compared to the other known toxic compositions.

For the purpose of practicing the invention, the predictions of toxicity of the test composition based on its molecular profiles in ES cells, tissue, etc, preferably EB cells does not have to be 100% accurate. To have a major positive impact on the efficiency and costs of drug development, one only has to modestly increase the probability that the less toxic and thus more successful drug candidates are, for example, on the top half of a prioritized list of new drug leads.

Since the cells and tissue obtained in accordance with the present invention more closely resemble the in vivo situation compared to conventional cell based assays, the results obtained by the assays of the present invention are expected to correlate to in vivo teratogenicity or embryotoxicity of the tested compounds as well. The simple read-out of the reporter molecule assay is easier measured and interpreted than determining the influence of a substance by monitoring the embryonic and post-natal development of a test animal in all aspects and in different stages of development. This requires a number of animals to be sacrificed at different ages, whereas by employing the reporter molecule secreted in an developmental-specific manner, samples of body fluids from the animal can be obtained repeatedly and allowing the monitoring over an extended time period up to the life-time of the animal. This also reduces the variability of test results.

Several test substances can be combined and either added simultaneously or sequentially to gain information about possible enhancing or quenching effects. Thus a further aspect of the invention relates to the method described previously, wherein said contacting step further includes contacting said test sample or animal with at least one second test substance in the presence of said first test substance. Two or more substances tested in combination will provide information about their interaction in general.

A preferred embodiment of the methods according to the present invention involves adding a compound known to activate or inhibit differentiation process to the culture medium or animal, particularly if this test substance is a therapeutic agent or a mixture thereof. This screening may be done, for example, either because the compound is known to have a effect on the differentiation of certain cell types and is tested to determine its potential for guiding the differentiation of other cell types, or because a compound designed to have effects elsewhere may have unintended side effects. The last aspect applies particularly to therapeutic agents.

Both aspects of drug screening can be adapted to compound library screening methods as well as screening industrial chemicals from the shelf. Thus another embodiment of the invention encompasses the method described above, wherein preferably in a first screen said test substance is comprised in and subjected as a collection of test substances. Such a collection of test substances, usually a combinatorial compound library, can have a diversity of about 10^3

to about 10^5 and is successively reduced in running the method several times until the diversity has been reduced to down to a desired factor, preferably down to one compound or particular class of compounds. Optionally the method of the present invention can combined with other screening methods known in the art. Methods for the generation and use of peptidomimetic combinatorial libraries are described in the prior art, for example in Ostresh, Methods in Enzymology 267 (1996), 220-234 and Dorner, Bioorg. Med. Chem. 4 (1996), 709-715. Drug discovery by dynamic combinatorial libraries is described, for example, in Nat. Rev. Drug Discov. 1 (2002), 26-36 and Drug Discov. Today 7 (2002), 117-125. Compounds identified using the method of the invention can be further evaluated in single experiments employing other methods of the present invention.

Compounds and candidate agents are obtained from a wide variety of sources including libraries of synthetic or natural compounds, or as mentioned industrial chemicals; see supra. For example, numerous means are available for random and directed synthesis of a wide variety of organic compounds and biomolecules, including expression of randomized oligonucleotides and oligopeptides. Alternatively, libraries of natural compounds in the form of bacterial, fungal, plant and animal extracts are available or readily produced. For example, inhibition of tumor-induced angiogenesis and matrix-metalloproteinase expression in confrontation cultures of embryoid bodies and tumor spheroids by plant ingredients used in traditional chinese medicine has been described by Wartenberg et al. in Lab. Invest. 83 (2003), 87-98.

A preferred embodiment of the screening method is performed on an array as provided by the invention

25

The assay methods of the present invention can be in a conventional laboratory format or adapted for high throughput screening. The term "high throughput screening" (HTS) refers to an assay design that allows easy analysis of multiple samples simultaneously, and capacity for robotic manipulation. Another desired feature of high throughput assays is an assay design that is optimized to reduce reagent usage, or minimize the number of manipulations in order to achieve the analysis desired. Examples of assay formats include 96-well, 384-well or more-well plates, levitating droplets, and "lab on a chip" microchannel chips used for liquid handling experiments. It is well known by those in the art that as miniaturization of plastic

30

- molds and liquid handling devices are advanced, or as improved assay devices are designed, that greater numbers of samples may be performed using the design of the present invention. In the method of the invention, said one or more cells or tissues are preferably contained in a container, for example in a well of a microtiter plate, which may be a 24, 96, 384 or 1586 well plate. Alternatively, the cells can be introduced into a microfluidics device, such as those provided by Caliper (Newton, MA, USA). In another preferred embodiment, the method of the present invention comprises taking 3, 4, 5, 7, 10 or more measurements, optionally at different positions within the container.
- 10 The methods according to the invention can also being performed using cells that are genetically engineered to (over) express or inhibit the expression of a target gene. Examples for genetically altered stem cells are described in WO03/014326 which provides mammalian stem cells expressing the VEGF receptor VEGFR-1 or WO95/06409 disclosing human hematopoietic stem cells expressing recombinant DNA constructs encoding a recombinant
- 15 molecule containing a signal transducing region and an antigenic specificity region. Alternatively to using already altered stem cells, said target gene can be transfected at the same time as the construct of the present invention conferring the expression of the secreted reporter molecule.
- 20 HTS, combinatorial library screening as well as conventional assay formats can also be employed to identify novel compounds which have an effect on differentiation. Novel as well as compounds previously known to modulate cell differentiation can be used to manufacture a drug for this purpose. Thus a preferred embodiment of the present invention is a method of obtaining and manufacturing a drug which promotes or inhibits formation of specific cell
- 25 types comprising the steps of the methods described above, wherein an enhanced or reduced level or activity of the reporter gene product is indicative for the drug.
- Compounds of interest encompass numerous chemical classes, though typically they are organic molecules. Candidate agents comprise functional groups necessary for structural interaction with proteins, particularly hydrogen bonding, and typically include at least an
- 30 amine, carbonyl, hydroxyl or carboxyl group, preferably at least two of the functional chemical groups. The candidate agents often comprise cyclical carbon or heterocyclic structures and/or aromatic or polyaromatic structures substituted with one or more of the above functional groups. Candidate agents are also found among biomolecules including

peptides, saccharides, fatty acids, steroids, purines, pyrimidines, derivatives, structural analogs or combinations thereof.

Additionally, natural or synthetically produced libraries and compounds are readily modified through conventional chemical, physical and biochemical means, and may be used to produce combinatorial libraries. Known pharmacological agents may be subjected to directed or random chemical modifications, such as acylation, alkylation, esterification, amidification, etc. to produce structural analogs.

The compounds may also be included in a sample including fluids to which additional components have been added, for example components that affect the ionic strength, pH, total protein concentration, etc. In addition, the samples may be treated to achieve at least partial fractionation or concentration. Biological samples may be stored if care is taken to reduce degradation of the compound, e.g. under nitrogen, frozen, or a combination thereof. The volume of sample used is sufficient to allow for measurable detection, usually from about 0.1 μ l to 1 ml of a biological sample is sufficient.

Differentiation processes do not only play a major role in embryonic development but are also important for an adult organism since the continuous replacement of senescent or cells damaged by injury or disease is achieved by precursor cells differentiating into the cells needed. An important aspect of the present invention is thus a method of manufacturing an agent which supports wound healing and/or healing of damaged tissue comprising the steps described herein, wherein an enhanced level or activity of the reporter gene product is indicative for said agent.

Compounds identified, isolated and/or produced by the methods described above can also be used as lead compounds in drug discovery and preparation of drugs or prodrugs. This usually involves modifying the lead compound or a derivative thereof or an isolated compound as described hereinbefore such as modifying said substance to alter, eliminate and/or derivatize a portion thereof suspected causing toxicity, increasing bioavailability, solubility and/or half-life. Accordingly, a preferred embodiment of the present invention encompasses the methods described previously and further modifying said substance to alter, eliminate and/or derivatize a portion thereof suspected causing toxicity, increasing bioavailability, solubility and/or half-life. Methods for clinical compound discovery comprises for example ultrahigh-throughput screening (Sundberg, Curr. Opin. Biotechnol. 11 (2000), 47-53) for lead identification, and

structure-based drug design (Verlinde and Hol, *Structure* 2 (1994), 577-587) and combinatorial chemistry (Salemme et al., *Structure* 15 (1997), 319-324) for lead optimization. The various steps recited above are generally known in the art. For example, computer programs for implementing these techniques are available; e.g., Rein, *Computer-Assisted Modeling of Receptor-Ligand Interactions* (Alan Liss, New York, 1989). Methods for the preparation of chemical derivatives and analogues are well known to those skilled in the art and are described in, for example, Beilstein, *Handbook of Organic Chemistry*, Springer edition New York Inc., 175 Fifth Avenue, New York, N.Y. 10010 U.S.A. and *Organic Synthesis*, Wiley, New York, USA. Furthermore, peptidomimetics and/or computer aided design of appropriate derivatives and analogues can be used, for example, according to the methods described above. Methods for the lead generation in drug discovery also include using proteins and detection methods such as mass spectrometry (Cheng et al. *J. Am. Chem. Soc.* 117 (1995), 8859-8860) and some nuclear magnetic resonance (NMR) methods (Fejzo et al., *Chem. Biol.* 6 (1999), 755-769; Lin et al., *J. Org. Chem.* 62 (1997), 8930-8931). They may also include or rely on quantitative structure-action relationship (QSAR) analyses (Kubinyi, *J. Med. Chem.* 41 (1993), 2553-2564, Kubinyi, *Pharm. Unserer Zeit* 23 (1994), 281-290) combinatorial biochemistry, classical chemistry and others; see, for example, Holzgrabe and Bechtold, *Pharm. Acta Helv.* 74 (2000), 149-155.

In one embodiment, the method of the invention further comprises mixing the substance isolated or modified with a pharmaceutically acceptable carrier. Examples of carriers and methods of formulation may be found in Remington's *Pharmaceutical Sciences*.

Substances are metabolized after their in vivo administration in order to be eliminated either by excretion or by metabolism to one or more active or inactive metabolites (Meyer, *J. Pharmacokinet. Biopharm.* 24 (1996), 449-459). Thus, rather than using the actual compound or drug identified and obtained in accordance with the methods of the present invention a corresponding formulation as a pro-drug can be used which is converted into its active form in the patient by his/her metabolism. Precautionary measures that may be taken for the application of pro-drugs and drugs are described in the literature; see, for review, Ozama, *J. Toxicol. Sci.* 21 (1996), 323-329.

The present invention also relates to a kit containing specific reagents such as those described hereinbefore useful for conducting any one of the above described methods of the present

invention, containing for example the reporter gene construct described hereinbefore, multi- or pluripotent cells, standard compounds, and optionally culture medium, recombinant nucleic acid molecules, detection means for the reporter molecule, in particular for SEAP such as EDTA, L-Homoarginine, NBT/BCIP, fixative solution, incubation buffer, PBS, etc. Such a
5 kit would typically comprise a compartmentalized carrier suitable to hold in close confinement at least one container. The carrier would further comprise reagents useful for performing said methods. The carrier may also contain a means for detection such as labeled enzyme substrates or the like.

10 Yet another aspect of the present invention relates to a method of conducting a drug discovery business, comprising:

- providing one or more assay systems as described herein for identifying a modulator of cell differentiation; and/or
- conducting therapeutic profiling of modulators identified in the previous step, or further
15 analogs thereof, for efficacy and toxicity in animals according to the invention; and
- formulating a pharmaceutical preparation including one or more modulators identified in the previous step as having an acceptable therapeutic profile.

Utilizing the methods described above, the identity of a modulator of differentiation or a
20 derivative thereof are determined. Agents are identified by their ability to alter the level of secreted reporter molecule. For suitable lead compounds that are identified, further therapeutic profiling of the agent, or analogs thereof, can be carried out for assessing efficacy and toxicity in animals. Those compounds having therapeutic profiles after animal testing can be formulated into pharmaceutical preparations for use in humans or for veterinary uses. The
25 subject business method can include an additional step of establishing a distribution system for distributing the pharmaceutical preparation for sale, and may optionally include establishing a sales group for marketing the pharmaceutical preparation.

Instead of developing the identified cell differentiation modulating agents in house, further
30 drug development can also be achieved by a different company. Thus a further aspect of the present invention relates to a method of conducting a target discovery business comprising:

- providing one or more assay systems described herein for identifying modulators of cell differentiation;

- (optionally) conducting therapeutic profiling of modulators identified in the previous step for efficacy and toxicity in animals according to the invention; and
 - licensing, to a third party, the rights for further drug development and/or sales for modulators identified in the first step, or analogs thereof.
- 5 For suitable lead compounds that have been identified, further profiling of the agent, or further analogs thereof, can be carried out for assessing efficacy and toxicity in animals, depending on the modalities of the agreement with the respective third party. Further development of those compounds for use in humans or for veterinary uses will then be conducted by the third party. The subject business method will usually involve either the sale
- 10 or licensing of the rights to develop said compound but may also be conducted as a service, offered to drug developing companies for a fee.

The present invention also relates to modulators of cell differentiation identified according to the methods described above as well as to pharmaceutical compositions for use in the

15 modulation of cell differentiation comprising such a modulator.

The modulator according to the invention can be combined with suitable diluents or carriers, preferably those which are pharmaceutically acceptable. Examples of such carriers, diluents and methods of formulation may be found in Remington's Pharmaceutical Sciences. To form

20 a pharmaceutically acceptable composition suitable for effective administration, such compositions will contain an effective amount of the modulator. Carriers or diluents are usually sterile and non-toxic, and defined as vehicles commonly used to formulate pharmaceutical compositions for animal or human administration. The diluent is selected so as not to affect the biological activity of the combination. Examples of such diluents are

25 distilled water, physiological saline, Ringer's solutions, dextrose solution, and Hank's solution. In addition, the pharmaceutical composition or formulation may also include other carriers, adjuvants, or nontoxic, nontherapeutic, nonimmunogenic stabilizers and the like. A therapeutically effective dose refers to that amount of modulator which is sufficient to achieve the desired effect on differentiation of target cells.

30 Further examples of suitable pharmaceutical carriers are well known in the art and include phosphate buffered saline solutions, emulsions, such as oil/water emulsions, various types of wetting agents, sterile solutions etc. Compositions comprising such carriers can be formulated by well known conventional methods. Accordingly, the present invention also provides a method of making a pharmaceutical composition for use in modulating cell differentiation

comprising mixing a modulator of cell differentiation identified according to a method of the invention with a suitable diluent or carrier.

The use of a reporter gene construct, a cell, a cell aggregate, tissue, organ, implant or transplant, a non-human animal, a composition, an array or the apparatus defined herein in drug discovery or pharmacokinetic or pharmacological profiling is also preferred embodiment of the present invention. Hence, the means and methods of the present invention described herein-before can be used in a variety of applications including but not limited to "loss of function" assays with ES cells containing homozygous mutations of specific genes, "gain of function" assays with ES cells overexpressing exogenous genes, developmental analysis of teratogenic/embryotoxic compounds in vitro, pharmacological assays and the establishment of model systems for pathological cell functions, and application of differentiation and growth factors for induction of selectively differentiated cells which, can be used as a source for tissue grafts; see for review, e.g., Guan et al., *Altex* 16 (1999), 135-141.

As described in the examples, for the expression of the secreted human placental alkaline phosphatase protein (SEAP), a particular promoter has been used, i.e. the promoter region of the mouse alpha myosin heavy chain gene; see NCBI GenBank Accession No. #U71441 (SEQ ID NO: 1). In accordance with the present invention, it could be shown that this promoter region is particularly suited for cell type specific expression of a reporter gene, especially for the purposes of the present invention. In addition, further experiments performed in accordance with the present invention surprisingly revealed that the mouse ventricular myosin regulatory light chain gene promoter (see NCBI GenBank Accession No. #AF302688; SEQ ID NO: 2) is particularly suited for the expression of the reporter gene as well. Thus, the present invention also relates to a recombinant DNA molecule as described hereinbefore in a cell-type specific reporter system, comprising the promoter of the mouse alpha myosin heavy chain gene and/or of the mouse ventricular myosin regulatory light chain gene, in particular the sequence depicted in NCBI GenBank Accession No. #U71441 (SEQ ID NO: 1) and #AF302688 (SEQ ID NO: 2), respectively, or a fragment thereof. Typically, the recombinant DNA molecule is a vector and comprises other functional DNA sequences such as ori, bacterial selectable marker, polyadenylation signal, multiple cloning sites, and the like. Hence, such vector may also be used to express specifically for examples growth factors and the like or target cDNAs in order to elucidate their function during cell development. Thus, the present invention involves the use of the mentioned promoter regions for the

expression of a DNA sequence encoding a reporter protein or selectable marker; see supra. In a particular, the present invention relates to the expression vector described in Example 1 comprising the nucleotide sequence depicted in SEQ ID NO. 3 as well as to a corresponding vector, wherein the mouse alpha myosin heavy chain gene promoter has been replaced by the mouse ventricular myosin regulatory light chain gene promoter, and their use in the afore-described embodiments.

These and other embodiments are disclosed and encompassed by the description and examples of the present invention. Further literature concerning any one of the materials, methods, uses and compounds to be employed in accordance with the present invention may be retrieved from public libraries and databases, using for example electronic devices. For example the public database "Medline" may be utilized, which is hosted by the National Center for Biotechnology Information and/or the National Library of Medicine at the National Institutes of Health. Further databases and web addresses, such as those of the European Bioinformatics Institute (EBI), which is part of the European Molecular Biology Laboratory (EMBL) are known to the person skilled in the art and can also be obtained using internet search engines. An overview of patent information in biotechnology and a survey of relevant sources of patent information useful for retrospective searching and for current awareness is given in Berks, TIBTECH 12 (1994), 352-364.

The above disclosure generally describes the present invention. A more complete understanding can be obtained by reference to the following specific examples and figure which are provided herein for purposes of illustration only and are not intended to limit the scope of the invention. The contents of all cited references (including literature references, issued patents, published patent applications as cited throughout this application and manufacturer's specifications, instructions, etc) are hereby expressly incorporated by reference; however, there is no admission that any document cited is indeed prior art as to the present invention.

The practice of the present invention will employ, unless otherwise indicated, conventional techniques of cell biology, cell culture, molecular biology, transgenic biology, microbiology, recombinant DNA, and immunology, which are within the skill of the art. For further elaboration of general techniques concerning stem cell technology, the practitioner can refer to standard textbooks and reviews, for example Teratocarcinomas and embryonic

stem cells: A practical approach (E. J. Robertson, ed., IRL Press Ltd. 1987); Guide to Techniques in Mouse Development (P. M. Wasserman et al., eds., Academic Press 1993); Embryonic Stem Cell Differentiation in Vitro (Wiles, Meth. Enzymol. 225 (1993), 900.); Properties and uses of Embryonic Stem Cells: Prospects for Application to Human Biology and Gene Therapy (Rathjen et al., Reprod. Fertil. Dev. 10 (1998), 31.). Differentiation of stem cells is reviewed in Robertson, Meth. Cell Biol. 75 (1997), 173; and Pedersen, Reprod. Fertil. Dev. 10 (1998), 31. Besides the sources for stem cells described already above further references are provided; see Evans and Kaufman, Nature 292 (1981), 154-156; Handyside et al., Roux's Arch. Dev. Biol., 196 (1987), 185-190; Flechon et al., J. Reprod. Fertil. Abstract Series 6 (1990), 25; Doetschman et al., Dev. Biol. 127 (1988), 224-227; Evans et al., Theriogenology 33 (1990), 125-128; Notarianni et al., J. Reprod. Fertil. Suppl., 43 (1991), 255-260; Giles et al., Biol. Reprod. 44 (Suppl. 1) (1991), 57; Strelchenko et al., Theriogenology 35 (1991), 274; Sukoyan et al., Mol. Reprod. Dev. 93 (1992), 418-431; Iannaccone et al., Dev. Biol. 163 (1994), 288-292.

15

Methods in molecular genetics and genetic engineering are described generally in the current editions of Molecular Cloning: A Laboratory Manual, (Sambrook et al., (1989) Molecular Cloning: A Laboratory Manual, 2nd ed., Cold Spring Harbor Laboratory Press); DNA Cloning, Volumes I and II (D. N. Glover ed., 1985); Oligonucleotide Synthesis (M. J. Gait ed., 1984); Nucleic Acid Hybridization (B. D. Hames & S. J. Higgins eds. 1984); Transcription And Translation (B. D. Hames & S. J. Higgins eds. 1984); Culture Of Animal Cells (R. I. Freshney, Alan R. Liss, Inc., 1987); Gene Transfer Vectors for Mammalian Cells (Miller & Calos, eds.); Current Protocols in Molecular Biology and Short Protocols in Molecular Biology, 3rd Edition (F. M. Ausubel et al., eds.); and Recombinant DNA Methodology (R. Wu ed., Academic Press). Gene Transfer Vectors For Mammalian Cells (J. H. Miller and M. P. Calos eds., 1987, Cold Spring Harbor Laboratory); Methods In Enzymology, Vols. 154 and 155 (Wu et al. eds.); Immobilized Cells And Enzymes (IRL Press, 1986); B. Perbal, A Practical Guide To Molecular Cloning (1984); the treatise, Methods In Enzymology (Academic Press, Inc., N.Y.); Immunochemical Methods In Cell And Molecular Biology (Mayer and Walker, eds., Academic Press, London, 1987); Handbook Of Experimental Immunology, Volumes I-IV (D. M. Weir and C. C. Blackwell, eds., 1986). Reagents, cloning vectors, and kits for genetic manipulation referred to in this disclosure are available from commercial vendors such as BioRad, Stratagene, Invitrogen, and Clontech. General techniques in cell culture and media collection are outlined in Large Scale

30

Mammalian Cell Culture (Hu et al., Curr. Opin. Biotechnol. 8 (1997), 148); Serum-free Media (Kitano, Biotechnology 17 (1991), 73); Large Scale Mammalian Cell Culture (Curr. Opin. Biotechnol. 2 (1991), 375); and Suspension Culture of Mammalian Cells (Birch et al., Bioprocess Technol. 19 (1990), 251). Other observations about the media and their impact on
5 the culture environment have been made by Marshall McLuhan and Fred Allen.

EXAMPLES

Example 1: Generation of a vector for cardiac-specific SEAP expression

Using standard cloning procedures, a SEAP expression vector was obtained in the following way. A 5.4 kb fragment containing the promoter region of the α -mouse myosin heavy chain gene (Genbank No. U71441) was inserted into the polylinker of the pEGFP-1 vector (Clontech). The α -myosin heavy chain gene is specifically expressed by cardiac myocytes in differentiating ES cells (Gulick et al., J. Biol. Chem. 266 (1991), 9180-91855; Palermo et al., Cell. Mol. Biol. Res. 41 (1995), 501-519; Morkin, Microsc. Res. Tech. 50 (2000), 522-531; Boheler et al., Circ. Res. 91 (2002), 189-201). Subsequently, the EGFP encoding sequence was removed and replaced by a SEAP encoding sequence (Berger et al., Gene 66 (1988), 1-10).

The resulting vector, designated α -MHC-SEAP, thus contains the SEAP gene under the transcriptional control of a heart-specific promoter. Further features of the vector include the presence of a Kozak consensus sequence to allow efficient translation of the SEAP mRNA, the presence of SV40 polyadenylation signals directing proper processing of the SEAP mRNA, the presence of a neomycin resistance cassette consisting of the SV40 early promoter, the neomycin/kanamycin resistance gene of Tn5 and polyadenylation signals from the Herpes simplex virus thymidine kinase gene to allow selection of stably transfected ES cells, and the presence of a bacterial promoter upstream of this cassette to confer kanamycin resistance in *E. coli*.

Example 2: Generation of ES cell transfectants

The α -MHC-SEAP vector was linearized by restriction endonuclease digestion and introduced into D3 mouse embryonic stem (ES) cells (American Type Culture Collection) by electroporation. Following neomycin selection, single ES cell colonies were isolated and expanded. ES cell culture and transfection as well as selection of stable transfectants was done using standard procedures (Torres and Kühn, Laboratory protocols for conditional gene targeting. Oxford University Press, (1997) New York).

Example 3: Differentiation of ES cells into cardiac cells in the presence of retinoic acid

The presence of excessive retinoic acid has striking effects on early heart development *in vivo* (Osmond et al., Development 113 (1991), 1405-1417; Yutzey et al., Development 120 (1994), 871-883; Dickman and Smith, Dev. Dyn. 206 (1996), 39-48; Drysdale et al., Dev. Biol. 188 (1997), 205-215). Furthermore, all-*trans* retinoic acid (CAS No. 302-79-4) was shown to inhibit the generation of cardiomyocytes from ES cells *in vitro* (Seiler et al. ALTEX 19 Suppl. 1 (2002), 55-63). Therefore, ES cells stably transfected with the α -MHC-SEAP vector were treated with retinoic acid at different concentrations and subjected to a differentiation protocol that resulted in the generation of cardiac cells. If secreted reporter, here SEAP, activity can indeed be used to quantify ES cell-derived cardiomyocytes, the amount of SEAP activity in culture supernatants should correlate inversely with the concentrations of retinoic acid present in the cultures.

All incubations of cell cultures were performed at 37°C in a humidified atmosphere containing 5% CO₂.

ES cell aggregation

ES cells stably transfected with the α -MHC-SEAP vector (see Example 2) were trypsinised to obtain a single cell suspension. Cells were collected by centrifugation (800 g for 5 min) and resuspended in Iscove's Modified Dulbecco's Medium (IMDM, Invitrogen) supplemented with 20% (v/v) fetal bovine serum (FBS, Invitrogen) at a density of 2×10^6 cells/ml. To aggregate ES cells, 4 ml of this suspension were incubated on a 6 cm bacteriological dish (Greiner) on a rocking platform (GFL 3006) with agitation (50 rpm). After 6 hours, 2 ml of that culture were added to 13 ml IMDM containing 20% FBS in a T25 flask (Falcon). The flask was incubated with rocking (50 rpm; GFL 3006) for 18 hours. During this procedure ES cell aggregates (embryoid bodies; EBs) were formed.

ES cell differentiation in the presence of retinoic acid

All-*trans* retinoic acid (Sigma) was dissolved in dimethyl sulfoxide (DMSO, Sigma) to obtain 10^{-4} M, 10^{-5} M, 10^{-6} M, 10^{-7} M, 10^{-8} M, and 10^{-9} M stock solutions. In 10 cm petri dishes (bacteriological grade, Greiner) medium was prepared consisting of IMDM supplemented with 20% FBS and retinoic acid added from above stocks to yield final concentrations of 10^{-7} M, 10^{-8} M, 10^{-9} M, 10^{-10} M, 10^{-11} M, and 10^{-12} M retinoic acid, respectively. One dish received DMSO (final concentration 0.1% (v/v)) without retinoic acid (solvent control). Each

dish contained 10 ml of medium. 70 EBs were transferred into each petri dish and incubated for 4 days.

- Subsequently, EBs were transferred to 6-well plates (tissue culture grade; Becton-Dickinson).
 5 Each well contained 3 ml IMDM with 20% FBS and retinoic acid and DMSO as described above for the culture in 10 cm dishes. 2 wells were set up for each concentration of retinoic acid and 2 wells as solvent controls. To each well 20 EBs were transferred from the respective 10 cm dish. Plates were incubated for 7 days. Subsequently, culture medium was exchanged with 2 ml knockout™ D-MEM (Invitrogen) supplemented with 20% (v/v) knockout™ SR
 10 (Invitrogen). Samples of cell culture supernatants were collected after incubation of EBs for further two days.

Determination of SEAP activity

- Samples were assayed for SEAP activity using the Great EscAPE SEAP Fluorescence
 15 Detection Kit (Becton-Dickinson) following the manufacturer's instructions. Fluorescence was measured using a microplate reader (Safire, Tecan). Background activity was determined by assaying a sample of fresh medium (knockout™ D-MEM supplemented with 20% (v/v) knockout™ SR).
 20 Results are shown in Fig. 1. Concentrations of retinoic acid below 10^{-9} M had no influence on SEAP activity as compared to controls. SEAP activity decreased sharply at 10^{-8} M retinoic acid. At higher concentrations no activity was observed.

- Thus, a concentration-dependent effect of retinoic acid on SEAP activity in culture
 25 supernatants is clearly observed. Moreover, these data are consistent with the concentration-dependent inhibition of cardiomyocyte generation by retinoic acid described previously (Seiler et al., 2002, supra). Therefore, the results indicate the suitability of the reporter system presented here to quantify ES cell derived cardiomyocytes and its usefulness for the monitoring of the differentiation process of ES cells

SEQUENCE LISTING

EPO-Maria
34
08. Juli 2003

5 <110> Axiogenesis AG

10 <120> Secreted Proteins As Markers For Cell Differentiation

15 <130> AX02A03/P-EP

20 <160> 4

25 <210> 1

 <211> 5443

 <212> DNA

30 <213> Mus musculus

35 <220>

 <221> promoter

 <222> (1)..(5443)

40 <223> alpha myosin heavy chain gene, promoter region

45 <220>

 <221> promoter

 <222> (2)..(5442)

50 <223> promoter region contained in SEQ ID NO 3

55 <220>

 <221> 5'UTR

 <222> (4369)..(4389)

60 <223> exon1

<220>

5 <221> 5'UTR

<222> (5071)..(5139)

<223> exon 2

10

<220>

15 <221> 5'UTR

<222> (5431)..(5443)

<223> exon 3

20

<400> 1

25	ggatcctgca aggtcacaca agggctctcca cccaccagggt gccctagtct caatttcagt	60
	ttccatgcct tgttctcaca atgctggcct cccagagct aatttggact ttgtttttat	120
	ttcaaaagggt cctgaatgag gagtagatct tgtgctaccc agctctaagg gtgcccggtga	180
30	agccctcaga cctggagcct ttgcaacagc cctttagggtg gaagcagaat aaagcaattt	240
	tccttaaagc caaaatcctg cctctagact cttctttctct gacctcggtc cctgggctct	300
	aggggtgggga ggtggggcctt ggaagaagaa ggtggggaag tggcaaaagc cgatccctag	360
35	ggccctgtga agttcggagc cttccctgta cagcactggc tcatagatcc tcctccagcc	420
	aaacatagca agaagtgata cctcctttgt gacttcccca ggcccagtac ctgtcagggt	480
40	gaaacaggat ttagagaagc ctctgaactc acctgaactc tgaagctcat ccaccaagca	540
	agcacctagg tgccactgct agttagtatc ctacgctgat aatatgcaga gctggggccac	600
	agaagtcctg ggggtgtagga actgaccagt gacttttcag tcggcaaagg tatgaccccc	660
45	tcagcagatg tagtaatgtc cccttagatc ccatcccagg caggtctcta agaggacatg	720
	ggatgagaga tgtagtcatg tggcattcca aacacagcta tcacagtgt cccttgcccc	780
50	ttccacttag ccaggaggac agtaacctta gcctatcttt cttoctcccc atcctoccag	840
	gacacacccc ctgggtctgca gtattcattt cttocttcac gtcccctctg tgacttccat	900
	ttgcaaggct tttgacctct gcagctgctg gaagatagag tttggcccta ggtgtggcaa	960
55	gccatctcaa gagaaagcag acaacagggg gaccagattt tggaaggatc aggaactaaa	1020
	tcactggcgg gcctgggggt agaaaaaaga gtgagtgagt ccgctccagc taagccaagc	1080
60	tagtccccga gatactctgc cacagctggg ctgctcgggg tagctttagg aatgtgggtc	1140
	tgaaagacaa tgggattgga agacatctct ttgagtctcc cctcaacccc acctacagac	1200

	acactcgtgt gtggccagac tcctgttcaa cagccctctg tgttctgacc actgagctag	1260
	gcaaccagag catgggccct gtgctgagga tgaagagttg gttaccaata gcaaaaacag	1320
5	caggggaggg agaacagaga acgaaataag gaaggaagaa ggaaaggcca gtcaatcaga	1380
	tgcagtcaga agagatggga agccaacaca cagcttgagc agaggaaaca gaaaaggag	1440
10	agattctggg cataaggagg ccacagaaag aagagcccag gcccccaag tctcctcttt	1500
	ataccctcat cccgtctccc aattaagccc actcttcttc ctagatcaga cctgagctgc	1560
	agcgaagaga cccgtaggga ggatcacact ggatgaagga gatgtgtgga gaagtccagg	1620
15	gcaacctaag agccagagcc taaaagagca agagataaag gtgcttcaaa ggtggccagg	1680
	ctgtgcacac agagggtcga ggactggtgg tagagcctca agataaggat gatgctcaga	1740
20	atgggcgggg ggggggattc tggggggggg agagagaagg tgagaaggag cctggaacag	1800
	agaatctgga agcgttgga acgataccat aaagggaaga acccaggcta cctttagatg	1860
	taaatcatga aagacaggga gaagggaagc tggagagagt agaaggaccc cggggcaaga	1920
25	catggaagca aggacaagcc aggttgagcg ctccgtgaaa tcagcctgct gaaggcagag	1980
	ccctggtatg agcaccagaa cagcagaggc taggggttaat gtcgagacag ggaacagaag	2040
30	gtagacacag gaacagacag agacggggga gccaggtaac aaaggaatgg tccttctcac	2100
	ctgtggccag agcgtccatc tgtgtccaca tactctagaa tgttcatcag actgcagggc	2160
	tggcttgga ggcagctgga aagagtatgt gagagccagg ggagacaagg gggcctagga	2220
35	aaggaagaag agggcaaacc aggccacaca agagggcaga gcccagaact gagttaactc	2280
	cttccttggt gcattctcca taggaggcag tgggaactct gtgaccacca tccccatga	2340
40	gccccacta cccataccaa gtttggcctg agtggcattc taggttcctt gaggacagag	2400
	cctggccttt gtctcttgga cctgacccaa gctgacccaa tgttctcagt accttatcat	2460
	gccctcaaga gcttgagaac caggcagtga catattaggc catgggctaa ccctggagct	2520
45	tgcacacagg agcctcaagt gacctccagg gacacagctg cagacagggt gcctttatcc	2580
	ccaaagagca accatttggc ataggtggct gcaaatggga atgcaagggt gaatcagggtc	2640
50	ccttcaagaa tactgcatgc aagacctaa acccctggag agaggggtat gtcctgccc	2700
	ccaccaccca taaggggagt gaactatcct agggggctgg cgacctggg gagacaccac	2760
	attactgaga gtgctgagcc cagaaaaact gaccgcctg tgcctgccc acctccacac	2820
55	tctagagcta tattgagagg tgacagtaga taggggtgga gctggtagca gggagagtgt	2880
	tcctgggtgt gaggtgtag gggaaagcca gagcagggga gtctggcttt gtctcctgaa	2940
60	cacaatgtct acttagttat aacaggcatg acctgctaaa gacccaacat ctacgacctc	3000
	tgaaaagaca gcagccctgg aggacagggg ttgtctctga gccttgggtg cttgatgggtg	3060
	ccacaaagga gggcatgagt gtgagtataa ggccccagga gcgttagaga agggcacttg	3120

	ggaaggggtc agtctgcaga gcccctatcc atggaatctg gagcctgggg ccaactggtg	3180
5	taaatctctg ggccctgccag gcattcaaag cagcacctgc atcctctggc agcctgggga	3240
	ggcggaaagg agcaaccccc cacttatacc ctttctccct cagccccagg attaacacct	3300
	ctggccttcc cccttccac ctcccatcag gagtggaggg ttgcagaggg agggtaaaaa	3360
10	cctacatgtc caaacatcat ggtgcacgat atatggatca gtatgtgtag aggcaagaaa	3420
	ggaaatctgc aggcttaact gggttaatgt gttaaagtctg tgtgcatgtg tgtgtgtctg	3480
15	actgaaaacg ggcattggtg tgcagctgtt cagttctgtg cgtgaggtta ccagactgca	3540
	ggtttgtgtg taaattgccc aaggcaaagt ggggtgaatcc cttccatggt ttaaagagat	3600
	tggatgatgg cctgcatctc aaggaccatg gaaaatagaa tggacactct atatgtgtct	3660
20	ctaagctaag gtagcaaggt ctttggagga cacctgtcta gagatgtggg caacagagac	3720
	tacagacagt atctgtacag agtaaggaga gagaggaggg ggtgtagaat tctcttacta	3780
25	tcaaagggaa actgagtcgt gcacctgcaa agtggatgct ctccctagac atcatgactt	3840
	tgtctctggg gagccagcac tgtggaactt cagggtctgag agagtaggag gctccctca	3900
	gcctgaagct atgcagatag ccagggttga aagggggaag ggagagcctg ggatgggagc	3960
30	ttgtgtgttg gaggcagggg acagatatta agcctggaag agaaggtgac ccttaccag	4020
	ttgttcaact cacccttcag attaaaaata actgaggtaa gggcctgggt aggggaggtg	4080
35	gtgtgagacg ctctgtctc tcctctatct gccatcggc cctttgggga ggaggaatgt	4140
	gccaaggac taaaaaagg ccatggagcc agaggggga gggcaacaga cctttcatgg	4200
	gcaaaccctg gggccctgct gtcctcctgt cacctccaga gccaagggat caaaggagga	4260
40	ggagccagga caggagggaa gtgggagggg ggggtccagc agaggactcc aaatttaggc	4320
	agcaggcata tgggatggga tataaagggg ctggagcact gagagctgtc agagatttct	4380
45	ccaaccaggy taagagggag tttcgggtgg gggctcttca cccacaccag acctctcccc	4440
	acctagaagg aaactgcctt tcctggaagt ggggttcagg ccggtcagag atctgacagg	4500
	gtggccttcc accagcctgg gaagtctca gtggcaggag gtttccacaa gaaacactgg	4560
50	atgcccttc ccttacgtg tcttctccat cttcctcctg gggatgctcc tcccgtctt	4620
	ggtttatctt ggctcttctg cttcagcaag atttgccctg tgetgtccac tccatcttct	4680
55	tctactgtct ccgtgccttg ccttgccttc ttgctgttcc ttcctttcca ccattttctc	4740
	acttcaacct ttctccctt ctcatctgta ttcactcttc cttccttctt tccttccctc	4800
	cttcccttct tccttccctt ctttctccct tccttccctt cttccttctt tccttccctt	4860
60	cttcccttct gtgtcagagt gctgagaatc acacctgggg tcccaccct tatgtaaaca	4920
	atcttccagt gagccacagc ttcagtgtct ctgggtgtct tcttaccttc ctacccccct	4980

ggcttgctct gttccatcct ggtcaggatc tctagattgg tctcccagcc tctgctactc 5040
 ctcttcctgc ctgttcctct ctctgtccag ctgcgccact gtggtgcctc gttccagctg 5100
 5 tgggtccacat tcttcaggat tctctgaaaa gttaaccagg tgagaatgtt tcccctgtag 5160
 acagcagatc acgattctcc cggagtcag gcttcagcc ctctctttct ctgcccagct 5220
 gcccggcact cttagcaaac ctcaggcacc cttacccac atagacctct gacagagaag 5280
 10 caggcacttt acatggagtc ctggtgggag agccataggc tacggtgtaa aagaggcagg 5340
 gaagtggtag tgtaggaaag tcaggacttc acatagaagc ctagccca cagaaatga 5400
 15 cagacagatc cctcctatct ccccataag agtttgagtc gac 5443

<210> 2

20 <211> 6298

<212> DNA

25 <213> Mus musculus

<220>

30 <221> promoter

<222> (1)..(5445)

35 <223> ventricular myosin regulatory light chain gene (AF302688)

<220>

40 <221> 5'UTR

<222> (5446)..(5486)

45 <223> exon 1

<220>

50 <221> exon

<222> (5487)..(5489)

55 <223> exon 1

<220>

60 <221> exon

<222> (6209)..(6298)

<223> exon 2

5 <400> 2
 ctcgagtgct gggattaaag gcatgcacca ccatgctcag cctgttttct tttttaagag 60
 ttaatccttt taattatgtg tatgtatgtc tttgcatgga tatgcccacg agagtgcattg 120
 10 tgccctggaa gaccagaagg gggtgacaaa cccctgcagt tttgagcctc ctggcctggg 180
 tgctgggaat caaacttaat ttctctagaa gagcagagtc ctcttaacac ccaagccctc 240
 15 ttgccagccc ttatcttctt tgggttggtg gtttttgcca aagctaggag ttccagcccc 300
 agtcttttat actctgccct catggcttct gacatctcca actctgccag gggctgactc 360
 ttatcttcag aaagcaattg ttactttgac cccagcagt gctgagccac caccagggc 420
 20 ctgaggctag aagaaaatgt accctccttc catggctcctc ttccagaaagc actgtgaacc 480
 caggtaaagg tagcaggctc aggttcagat gtagtgtcca cagcctgtgc aaaaaggacc 540
 cttcacttcc aagcttgccc cacctgatca gaccactctc cagatctggg ggaaacggta 600
 25 gcaggctctt caccccaaca ggagtcccca gggcacacaa gatattgcct ttgattttgt 660
 ccttatgaca accctgcgaa taaaatcgtc atcatgccc ctctacagat tagaatactg 720
 30 aagctgagag caaaaaccta cctgatgtca cactgctgct caatgactaa gcaacgaaaa 780
 tcaaaagtga atattccgc aggtccagt cttgcagatg acaaatgctc tgtcaccgtc 840
 acatttggtt ccatgttctg aaatcctctg tttcacctc cctagaacct cagatagtag 900
 35 ctaaatagct gaggccattc acctggccct ggctttctgc taaggaatat gtcagacatc 960
 ttccaggattc ttcaaacctg ttctgagtag aaatctctc aggcctggga gctgtttcct 1020
 40 ctccagcaga agccccatg tggcacaggc caactcccc acaaccagaa ccatcatcag 1080
 tcacaactgc acttcctctc atgccgaaga aactttttt ttccagtctg gatggatgct 1140
 ggaaccatcg gatcctgtaa ccgcttctc cactccagt acagacgtca ggacttgagg 1200
 45 ctccctccgt ctgctctctt ctgcactgca cttgaatttc cctgagggtt ggcatattgg 1260
 gtttttggtt ggcttggtt tttgcttgcc tagaggaata atcaatgctt aggcctttta 1320
 50 tagcaaaggc atttagtggg tcaaccaagc actcaggaga tagacaggag ggtcagaatt 1380
 caaggctcct ctttcttaca caatgagttc agggctcagc tgggctacac gagatccaat 1440
 ctaaaataaa caaaaaataa aataattgat atttcttctc tgtatattat agttatactg 1500
 55 tagactgttg gggatacatt ctccaagagt ctcaaagttc atctccactg agttaattaa 1560
 tggtgttact ataacacaac actagagtct ggttcgttta caaatacaca agattttatt 1620
 60 tgggctggag agatggctca gagctgactg ctcttccaaa ggtcctgagt tccaatccca 1680
 gcaaccacat ggtggctcga aaccatctgt aatgggatct gatgccttct tctagtgtgt 1740

	ctgaagacag	ctatggtgta	ctcacatata	taaaataaat	aaataaatct	taagagagag	1800
	atttgtttct	acaaaggcta	gatgctactg	agagtaaagg	gtttagaact	ccgttttcct	1860
5	cttcttacta	actcactaat	cccatcttgg	gggggggagg	gggaagcttc	atgacttcac	1920
	ctaacctcgt	tcacctctcc	tagaccttac	ctcctatcgc	taccaatctc	tgatttgggg	1980
10	atcaaatttt	caaccagga	gctttaggga	gacaaatgta	aacccagta	tccoctgact	2040
	cccggtgtac	actggatcct	gcacacagta	gggctccaat	aaatactgaa	agtccttaga	2100
	atgggggcaa	aatcattttt	gtgtttataa	atatgaccca	catgttctct	catcttttaa	2160
15	aattgtcaga	gtaatttctc	tctctctctc	tctctctctc	tctgtgtgtg	tgtgtgtgtc	2220
	tgtgtctgtg	tgtgtgtctg	tgtctgtgtc	tgtgtgtctg	tgtgtgtgta	ctcacatcca	2280
	agtgggtttg	ggtacctgtc	atgtggacat	gtatatgttg	gtagaagcca	gaagtcaacc	2340
20	ttgtgtgtca	tttctcagag	gtgatccatc	ttattttttg	agacagggtc	tctcactgag	2400
	gccggggact	tactgttttt	aggttacacc	tgctggccag	caagccagag	agaccgagag	2460
25	agacggtctg	tctccatctc	tccagcgaca	ggctttcaga	cccacaccac	catgcccagc	2520
	tttgatgtgg	gtctgggttt	tgaggagtag	aattcaggac	ctcttgctga	gccatctccc	2580
	cagccactga	acataattca	tatataatct	ggcttttggt	ccttttggtg	gtgcagggtg	2640
30	tgttttgttt	tgttctttgc	tgtgtattat	tctgtgcatt	taagcttttt	ttttttgtga	2700
	ctacctggga	gggacactga	acagccagaa	aggccaggcc	gaggcttcac	tgatggggat	2760
35	gtgagcccg	gaatgctggc	agctgccggg	agctggaaag	ggcaaaggaa	aggactgtct	2820
	ccgcatccac	ggtggacgca	gccctctgcc	cgatttctat	ttctggccac	cagctctgcc	2880
40	aggtagcgag	cttggtctgt	ctgagctgcc	tgggtttgcc	gttgattttt	cctactagca	2940
	tgggaaagcg	tgatcagctt	gctttgttct	caattgttcc	agaagctctg	ccggtcccct	3000
	ccaggactcc	tgagtctctg	ctccgtggta	ctcggggctg	gctctcaaag	ttctaggctg	3060
45	cagaaatctc	acaagcgcat	gagcttagag	tcccagggtg	gatgtccacc	aagagcccag	3120
	ggacaaagca	ttgacagctc	ctgtgtgcgc	cacgtctccc	cccaccccca	cccctacccc	3180
50	caggaactgt	gagaggagt	cagagcccct	ccccaggcc	ttccaacaa	ggactcctgg	3240
	aggacccttg	ggtttttaac	accaaataac	caaagtgttc	ccacgcagca	acacaaacca	3300
	gtcttctctc	atacagcacg	gtgggccaagt	ggaccatggg	gacagggttac	ctctgtgggc	3360
55	ccaggctcac	ggtaaactct	aacctcaatc	tgtagcctcc	cacagccatt	tgcgggtcac	3420
	cttgcttctc	agccaccgtg	tggcacttgg	caagtcacgt	gtgcctcaac	acaataagaa	3480
	gccaaaggaa	taggggcttt	gcttaactgg	tacagcagtg	tagcccaagc	tagccttgaa	3540
60	ctcactatgt	agccaaggac	gatcataaac	tcttgatcct	cccgttccag	agtcctgggt	3600
	gttgagataa	cagggtgtgg	tactcccta	cccttcttct	aatagcaatc	aatgtgtggc	3660

	cacatgtttg tgcctcacag attaaaacca tcttgacctg aggacgaaat gactaacagt	3720
	tgcctcctga aggttgacctg gatctcatct ttataatccc agcaatcaag gggagtgggg	3780
5	gatcaggagt tcaaagtcag ccagacctgg gctacatgag accctgtctt gaaaaatgga	3840
	ggaattaagc tgggcgtggg gccgcactcc tttaatccca gcacttggga ggtagaggca	3900
10	ggcggacttc tgagttcgag gccagacctgg tctacaaagt gagttccagg acagccacag	3960
	ctatacagag aaacctgtc tcgaaaaacc agaaagaaag aaagagagaa agaaagaaag	4020
	aaagaaagaa agaaagaaag aaagaaagga aggaaggaag gaaggaagga aaggaaggaa	4080
15	ggaaggaagg aaagaaagaa agaaagaaag aaagaaagac agacagaaag aaagggttagg	4140
	aaagaaagaa aggaaaagaa agaaagaaag aaagaaagaa agaaagaaag aaagaaagaa	4200
20	agacagaaag aaaggtttgg aaagaaagaa aggaaaagaa agaaagaaag agagaaagag	4260
	agaaagaaag aaagaaaaga aaagagagaa aagaaaagaa aagaaaagaa aagaaagaaa	4320
	agaaaaaaaa gaaaagaaaa gaaagaaaag aaaaggaagg aagaaaagaa aagaaaaatg	4380
25	gaggagttaa ccctatgttt cttttttttt tattcatcat tggtagaggct atcctcagct	4440
	acatatcaag ttcaagccag cctgggctac atgagaccct gcctcaaaaa agaaaaggag	4500
30	ccagtgtagc gacatactcc cgtcctccca gcacttggga gacagaggct actccactgc	4560
	tgtctccagc agccggcctg cctccctgag cctcattttt ttcataacat ggggacccaa	4620
	ctgctaaggt gaccttgctc ccatgggggtg actggagact tgagagtga gttggttatca	4680
35	tttgtccagt ctgtgaacaa atggcagcct ccaagggtgt tttgtgttca aaggaggaca	4740
	tgggacaggg agaggccagg gagaagagcc caccctcagg agtaggctgt ccccgtagag	4800
40	ctgggtgggg acaaaaagca gagaagcaga ggcagaggac aagcgtgggt gacatttgag	4860
	caaagatggg aatgtgccag aggctgccca agatgtgcat gtgcaaaggc cctgaggtgc	4920
	aagggtgcct ggatccagag ccaaaagctc aggetccctc ctctcttcc tctcttctc	4980
45	cctcttctc ctctttctc tctcctcct ctccctctc ccccttctc tctctctcc	5040
	cttctctc tctcttctc ccttctctc cctccctc ctctctctc tctctctct	5100
50	cctctctc ctctctctc tctctctct gctctctc ctctctctc tgggttactc	5160
	ttccccatta gacaatggca gggaaagagag cacaccccat catccccagg ccaggcccca	5220
	gccactgact ctttaacctt gaaggcattt ttgggtctca cgtgtccacc caggcgggtg	5280
55	gccgcctttg agcagctctt acttcagaag aacggcatgg agtgggggtt ggggggctta	5340
	ggtggcctcc gcctcaccta caactgcaa aagtggcat ggggttattt ttaaccccag	5400
60	gggagaggta ttattgttc cacagcaggg gcagaggcca gcaggctcct cgaactctcc	5460
	agagggtggca actggcctca gacacc atg gtgagtggtc agtaaaccct tgagaagaga	5519
	Met	

1

	cgaggggtgg ggagcagaga gataaccccc tcccaagcca agcaccccat ggaggagggg	5579
5	gggaggagga ggggaaggag gagggaagct ctcttacgag ccccttagcc ctagatggac	5639
	cagcaccttg caccctctga gggaccccaa tcagctcccc taaggagcca caaatagcag	5699
	ctcctcaagg aacttgcaaa aatcaatgag aaatgcgctt gggggatggg tgcccactac	5759
10	ctgatctcaa gaaatcagta acaccccacc cccaccccac ccagagcttg ccaaacggag	5819
	actgagagct ttttaagggc gaattgtaat tcttttccca attcaggtgg ccaggaagag	5879
15	gttttctggt tctctctttt gaattattccc cttgaaatat ttgtgcccgc ctcccagaac	5939
	aggtagcccc cagctgctag agactgcagc taaggggagc agagtgtacg tgtgtgtggc	5999
	tgtgtgatct agagaagtga ctcaccctct ctgagcctcc agtctcctta gtggagcaga	6059
20	ggagagcatt agataatgtt tggagggtttt ggggtatcat ttgcctcgc atgttgtctg	6119
	ggtaccagag actcactccc caggtgacag gtcctggccc aggtcctgat ccagaggctc	6179
25	cacagtgtct gatggatatt cctctccag gca cca aag aaa gcc aag aag cgg	6232
	Ala Pro Lys Lys Ala Lys Lys Arg	5
	ata gaa ggc ggg agc tcc aac gtg ttc tcc atg ttt gag cag acc cag	6280
30	Ile Glu Gly Gly Ser Ser Asn Val Phe Ser Met Phe Glu Gln Thr Gln	10 15 20 25
	atc cag gag ttc aag gaa	
35	Ile Gln Glu Phe Lys Glu	30
	<210> 3	
40	<211> 10475	
	<212> DNA	
	<213> Artificial Sequence	
45		
	<220>	
50	<223> composite vector	
	<220>	
	<221> promoter	
55	<222> (77)..(5517)	
	<223> 77-5517 Mus musculus alpha myosin heavy chain gene, promoter regi	
60	on (Acc. No. U71441)	

<220>
<221> CDS
5 <222> (5558)..(7117)
<223> SEAP

10 <220>
<221> rep_origin
15 <222> (9750)..(10393)
<223> pUC plasmid replication origin

20 <220>
<221> polyA_signal
25 <222> (9401)..(9406)
<223> Herpes simplex virus thymidine kinase polyadenylation signal

30 <220>
<221> polyA_signal
35 <222> (9414)..(9419)
<223> Herpes simplex virus thymidine kinase polyadenylation signal

40 <220>
<221> terminator
45 <222> (9163)..(9165)
<223> Kanamycin/neomycin resistance gene

50 <220>
<221> ATG
55 <222> (8371)..(8373)
<223> Kanamycin/neomycin resistance gene

60 <220>

<221> promoter
<222> (8242)..(8248)
5 <223> SV40 early promoter element

10 <220>
<221> repeat_region
<222> (8167)..(8230)
15 <223> SV40 early promoter

20 <220>
<221> enhancer
<222> (8018)..(8163)
25 <223> SV40 early promoter

30 <220>
<221> rep_origin
<222> (8187)..(8322)
35 <223> SV40 origin of replication

40 <220>
<221> -10_signal
<222> (7931)..(7936)
45 <223> Ampicillin resistance (b-lactamase) promoter

50 <220>
<221> -35_signal
<222> (7908)..(7913)
55 <223> Ampicillin resistance (b-lactamase) promoter

60 <220>
<221> polyA_signal

<222> (7294)..(7299)

<223> SV40 early mRNA polyadenylation signal

5

<220>

<221> polyA_signal

10

<222> (7323)..(7328)

<223> SV40 early mRNA polyadenylation signal

15

<220>

<221> mature protein

20

<222> (5609)..(7114)

<223> SEAP

25

<220>

<221> start codon

30

<222> (5558)..(5560)

<223> ATG of the Secreted alkaline phosphatase (SEAP) gene

35

<220>

<221> sig_peptide

40

<222> (5558)..(5608)

<223> SEAP

45

<400> 3	tagttattac tagcgctacc ggactcagat ctcgagctca agcttcgaat tctgcagtcg	60
50	acggtaccgc gggcccgatc ctgcaaggtc acacaagggc ctccacccac caggtgccct	120
	agtctcaatt tcagttttcca tgcccttggtc tcacaatgct ggccctcccca gagctaattt	180
	ggacttttgtt tttattttcaa aagggccctga atgaggagta gatcttggtc taccagctc	240
55	taaggggtgcc cgtgaagccc tcagacctgg agcctttgca acagcccttt aggtggaagc	300
	agaataaagc aattttcctt aaagccaaaa tctgcctct agactcttct tctctgacct	360
60	cggtcctctgg gctctagggc ggggaggtgg ggcttggaag aagaaggtgg ggaagtggca	420
	aaagccgatc cctagggccc tgtgaagttc ggagccttcc ctgtacagca ctggctcata	480

	gatcctcctc cagccaaaca tagcaagaag tgatacctcc tttgtgactt ccccaggccc	540
	agtacctgtc aggttgaaac aggatttaga gaagcctctg aactcacctg aactctgaag	600
5	ctcatccacc aagcaagcac ctaggtgcc a ctgctagtta gtatcctacg ctgataatat	660
	gcagagctgg gccacagaag tcctgggggtg taggaactga ccagtgactt ttcagtcggc	720
	aaaggtatga cccctcagc agatgtagta atgtccctt agatcccatc ccaggcaggt	780
10	ctctaagagg acatgggatg agagatgtag tcatgtggca ttccaaacac agctatccac	840
	agtgtccctt gccccttcca cttagccagg aggacagtaa ccttagccta tctttcttcc	900
15	tccccatcct ccaggacac acccctggt ctgcagtatt catttcttcc ttcacgtccc	960
	ctctgtgact tccatttgca aggcctttga cctctgcagc tgctggaaga tagagtttgg	1020
	ccctaggtgt ggcaagccat ctcaagagaa agcagacaac agggggacca gattttggaa	1080
20	ggatcaggaa ctaaactact ggcgggcctg ggggtagaaa aaagagttag tgagtccgct	1140
	ccagctaagc caagctagtc cccgagatac tctgccacag ctgggctgct cggggtagct	1200
25	ttaggaatgt ggggtctgaaa gacaatggga ttggaagaca tctctttgag tctcccctca	1260
	acccaccta cagacacact cgtgtgtggc cagactcctg ttcaacagcc ctctgtgttc	1320
	tgaccactga gctaggcaac cagagcatgg gccctgtgct gaggatgaag agttgggttac	1380
30	caatagcaaaa aacagcaggg gagggagaac agagaacgaa ataaggaagg aagaaggaaa	1440
	ggccagtcaa tcagatgcag tcagaagaga tgggaagcca acacacagct tgagcagagg	1500
35	aaacagaaaa gggagagatt ctgggcataa ggaggccaca gaaagaagag cccaggcccc	1560
	ccaagtctcc tctttatacc ctcatcccgt ctcccaatta agcccactct tcttcctaga	1620
	tcagacctga gctgcagcga agagaccctg agggaggatc aactggatg aaggagatgt	1680
40	gtggagaagt ccagggaac ctaagagcca gagcctaaaa gagcaagaga taaaggtgct	1740
	tcaaaggtgg ccaggctgtg cacacagagg gtgcaggact ggtggtagag cctcaagata	1800
45	aggatgatgc tcagaatggg cggggggggg gattctgggg gggggagaga gaaggtgaga	1860
	aggagcctgg aacagagaat ctggaagcgc tggaaacgat accataaagg gaagaaccca	1920
	ggctaccttt agatgtaaat catgaaagac agggagaagg gaagctggag agagtagaag	1980
50	gaccccgggg caagacatgg aagcaaggac aagccagggt gagcgtccg tgaaatcagc	2040
	ctgctgaagg cagagccctg gtatgagcac cagaacagca gaggctaggg ttaatgtcga	2100
55	gacagggaaac agaaggtaga cacaggaaca gacagagacg ggggagccag gtaacaaagg	2160
	aatggctcct ctccactgtg gccagagcgt ccatctgtgt ccacatactc tagaatgttc	2220
	atcagactgc agggctggct tgggaggcag ctggaaagag tatgtgagag ccaggggaga	2280
60	caagggggcc taggaaagga agaagagggc aaaccaggcc acacaagagg gcagagccca	2340
	gaactgagtt aactccttcc ttgttgcatc ttccatagga ggcagtggga actctgtgac	2400

	caccatcccc	catgagcccc	cactacccat	accaagtttg	gcctgagtg	cattctaggt	2460
5	tccttgagga	cagagcctgg	cctttgtctc	ttggacctga	cccaagctga	cccaatgttc	2520
	tcagtacctt	atcatgccct	caagagcttg	agaaccaggc	agtacatat	taggcatgg	2580
	gctaacctg	gagcttgac	acaggagcct	caagtgcct	ccagggacac	agctgcagac	2640
10	aggtggcctt	tatccccaaa	gagcaaccat	ttggcatagg	tggctgcaa	tgggaatgca	2700
	aggttgaatc	aggtcccttc	aagaatactg	catgcaagac	ctaagacccc	tggagagagg	2760
15	ggtatgctcc	tgccccacc	caccataagg	ggagtgaact	atcctagggg	gctggcgacc	2820
	ttggggagac	accacattac	tgagagtgtc	gagcccagaa	aaactgaccg	ccctgtgtcc	2880
	tgcccacctc	cacactctag	agctatattg	agaggtgaca	gtagataggg	tgggagctgg	2940
20	tagcaggag	agtgttctct	ggtgtgagg	tgtaggggaa	agccagagca	gggagtgctg	3000
	gctttgtctc	ctgaacacaa	tgtctactta	gttataacag	gcatgacctg	ctaaagaccc	3060
25	aacatctacg	acctctgaaa	agacagcagc	cctggaggag	aggggttgtc	tctgagcctt	3120
	gggtgcttga	tgggtgccaca	aaggagggca	tgagtgtgag	tataaggccc	caggagcgtt	3180
	agagaagggc	acttgggaag	gggtcagctc	gcagagcccc	tatccatgga	atctggagcc	3240
30	tggggccaac	tgggtgaaat	ctctgggcct	gccaggcatt	caaagcagca	cctgcatcct	3300
	ctggcagcct	ggggaggcgg	aaggagggca	cccccaactt	ataccctttc	tcctcagcc	3360
35	ccaggattaa	cacctctggc	cttccccctt	cccacctccc	atcaggagtg	gaggggtgca	3420
	gagggagggt	aaaaacctac	atgtccaaac	atcatgggtg	acgatatatg	gatcagtatg	3480
	tgtagaggca	agaaaggaaa	tctgcaggct	taactgggtt	aatgtgtaaa	gtctgtgtgc	3540
40	atgtgtgtgt	gtctgactga	aaacgggcat	ggctgtgcag	ctgttcagtt	ctgtgcgtga	3600
	ggttaccaga	ctgcaggttt	gtgtgtaaat	tgccaaggc	aaagtgggtg	aatcccttcc	3660
45	atggttttaa	gagattggat	gatggcctgc	atctcaagga	ccatggaaaa	tagaatggac	3720
	actctatatg	tgtctctaag	ctaaggtagc	aaggctcttg	gaggacacct	gtctagagat	3780
	gtgggcaaca	gagactacag	acagtatctg	tacagagtaa	ggagagagag	gagggggtgt	3840
50	agaattctct	tactatcaaa	gggaaactga	gtcgtgcacc	tgcaaagtgg	atgctctccc	3900
	tagacatcat	gactttgtct	ctggggagcc	agcactgtgg	aacttcaggt	ctgagagagt	3960
55	aggaggctcc	cctcagcctg	aagctatgca	gatagccagg	gttgaaagg	ggaagggaga	4020
	gcctgggatg	ggagcttgtg	tgttgaggc	aggggacaga	tattaagcct	ggaagagaag	4080
	gtgaccctta	cccagttgtt	caactcacc	ttcagattaa	aaataactga	ggtaagggcc	4140
60	tgggtagggg	aggtggtgtg	agacgtcctc	gtctctctc	tatctgcca	tcggcccttt	4200
	ggggaggagg	aatgtgcca	aggactaaaa	aaaggccatg	gagccagagg	ggcgagggca	4260

	acagaccttt catgggcaaa ccttggggcc ctgctgtcct cctgtcacct ccagagccaa	4320
	gggatcaaag gaggaggagc caggacagga gggaagtggg agggagggtc ccagcagagg	4380
5	actccaaatt taggcagcag gcatatggga tgggatataa aggggctgga gcactgagag	4440
	ctgtcagaga tttctccaac ccaggtaaga gggagtcttcg ggtgggggct cttcaccac	4500
10	accagacctc tccccaccta gaaggaaact gcctttcctg gaagtggggt tcaggccggt	4560
	cagagatctg acaggggtggc cttccaccag cctgggaagt tctcagtggc aggaggtttc	4620
	cacaagaaac actggatgcc ccttccctta cgctgtcttc tccatcttcc tcoctgggat	4680
15	gctcctcccc gtcttggttt atcttggtc ttcgtcttca gcaagatttg cctgtgtctg	4740
	tccactccat ctttctctac tgtctccgtg ccttgccttg ccttcttgcg tgctcttct	4800
20	ttccacccat ttctcacttc accttttctc ccttctctat ttgtattcat ccttcttcc	4860
	ttccttctct ccttcttcc ttcttctct ccttcttct tcccttctct ccttcttcc	4920
	ttccttctct ccttcttcc ttctgtgtc agagtgtga gaatcacacc tgggggtccc	4980
25	acccttatgt aaacaatctt ccagtggcc acagcttcag tgctgtggg tgctctctta	5040
	ccttctctac ccttgggtt gtctgttcc atctgtgtca ggaatctctag attggtctcc	5100
30	cagcctctgc tactcctctt cctgctgtt cctctctctg tccagctgag ccaactgtgg	5160
	gctctgttcc agctgtgtc cacattcttc aggattctct gaaaagttaa ccaggtgaga	5220
	atgtttcccc tgtagacagc agatcacgat tctcccggaa gtcaggcttc cagccctctc	5280
35	tttctctgcc cagctgcccg gcaactcttag caaacctcag gcacccttac cccacataga	5340
	cctctgacag agaagcaggc actttacatg gactcctggt gggagagcca taggctacgg	5400
40	tgtaaaagag gcaggggaagt ggtggtgtag gaaagtcagg acttcacata gaagcctagc	5460
	ccacaccaga aatgacagac agatccctcc tatctcccc ataagagttt gactcgaggg	5520
	atctaagcag aagcttcgaa tcggaattc gccacc atg ctg ctg ctg ctg ctg	5575
45	Met Leu Leu Leu Leu Leu 1 5	
	ctg ctg ggc ctg agg cta cag ctg tcc ctg ggc atc atc cca gtt gag	5623
	Leu Leu Gly Leu Arg Leu Gln Leu Ser Leu Gly Ile Ile Pro Val Glu	
50	10 15 20	
	gag gag aac ccg gac ttc tgg aac cgc gag gca gcc gag gcc ctg ggt	5671
	Glu Glu Asn Pro Asp Phe Trp Asn Arg Glu Ala Ala Glu Ala Leu Gly	
	25 30 35	
55	gcc gcc aag aag ctg cag cct gca cag aca gcc gcc aag aac ctg atc	5719
	Ala Ala Lys Lys Leu Gln Pro Ala Gln Thr Ala Ala Lys Asn Leu Ile	
	40 45 50	
60	atc ttc ctg ggc gat ggg atg ggg gtg tct acg gtg aca gct gcc agg	5767
	Ile Phe Leu Gly Asp Gly Met Gly Val Ser Thr Val Thr Ala Ala Arg	
	55 60 65 70	
	atc cta aaa ggg cag aag aag gac aaa ctg ggg cct gag ata ccc ctg	5815

	Ile	Leu	Lys	Gly	Gln	Lys	Lys	Asp	Lys	Leu	Gly	Pro	Glu	Ile	Pro	Leu	
					75					80					85		
5	gcc	atg	gac	cgc	ttc	cca	tat	gtg	gct	ctg	tcc	aag	aca	tac	aat	gta	5863
	Ala	Met	Asp	Arg	Phe	Pro	Tyr	Val	Ala	Leu	Ser	Lys	Thr	Tyr	Asn	Val	
				90					95					100			
10	gac	aaa	cat	gtg	cca	gac	agt	gga	gcc	aca	gcc	acg	gcc	tac	ctg	tgc	5911
	Asp	Lys	His	Val	Pro	Asp	Ser	Gly	Ala	Thr	Ala	Thr	Ala	Tyr	Leu	Cys	
			105					110					115				
15	ggg	gtc	aag	ggc	aac	ttc	cag	acc	att	ggc	ttg	agt	gca	gcc	gcc	cgc	5959
	Gly	Val	Lys	Gly	Asn	Phe	Gln	Thr	Ile	Gly	Leu	Ser	Ala	Ala	Ala	Arg	
		120					125					130					
20	ttt	aac	cag	tgc	aac	acg	aca	cgc	ggc	aac	gag	gtc	atc	tcc	gtg	atg	6007
	Phe	Asn	Gln	Cys	Asn	Thr	Thr	Arg	Gly	Asn	Glu	Val	Ile	Ser	Val	Met	
	135					140					145					150	
25	aat	cgg	gcc	aag	aaa	gca	ggg	aag	tca	gtg	gga	gtg	gta	acc	acc	aca	6055
	Asn	Arg	Ala	Lys	Lys	Ala	Gly	Lys	Ser	Val	Gly	Val	Val	Thr	Thr	Thr	
					155					160					165		
30	cga	gtg	cag	cac	gcc	tcg	cca	gcc	ggc	acc	tac	gcc	cac	acg	gtg	aac	6103
	Arg	Val	Gln	His	Ala	Ser	Pro	Ala	Gly	Thr	Tyr	Ala	His	Thr	Val	Asn	
				170					175					180			
35	cgc	aac	tgg	tac	tcg	gac	gcc	gac	gtg	cct	gcc	tcg	gcc	cgc	cag	gag	6151
	Arg	Asn	Trp	Tyr	Ser	Asp	Ala	Asp	Val	Pro	Ala	Ser	Ala	Arg	Gln	Glu	
			185					190					195				
40	ggg	tgc	cag	gac	atc	gct	acg	cag	ctc	atc	tcc	aac	atg	gac	att	gac	6199
	Gly	Cys	Gln	Asp	Ile	Ala	Thr	Gln	Leu	Ile	Ser	Asn	Met	Asp	Ile	Asp	
		200					205					210					
45	gtg	atc	cta	ggt	gga	ggc	cga	aag	tac	atg	ttt	cgc	atg	gga	acc	cca	6247
	Val	Ile	Leu	Gly	Gly	Gly	Arg	Lys	Tyr	Met	Phe	Arg	Met	Gly	Thr	Pro	
		215				220					225					230	
50	gac	cct	gag	tac	cca	gat	gac	tac	agc	caa	ggt	ggg	acc	agg	ctg	gac	6295
	Asp	Pro	Glu	Tyr	Pro	Asp	Asp	Tyr	Ser	Gln	Gly	Gly	Thr	Arg	Leu	Asp	
					235				240						245		
55	ggg	aag	aat	ctg	gtg	cag	gaa	tgg	ctg	gcg	aag	cgc	cag	ggt	gcc	cgg	6343
	Gly	Lys	Asn	Leu	Val	Gln	Glu	Trp	Leu	Ala	Lys	Arg	Gln	Gly	Ala	Arg	
				250				255					260				
60	tat	gtg	tgg	aac	cgc	act	gag	ctc	atg	cag	gct	tcc	ctg	gac	ccg	tct	6391
	Tyr	Val	Trp	Asn	Arg	Thr	Glu	Leu	Met	Gln	Ala	Ser	Leu	Asp	Pro	Ser	
			265				270						275				
65	gtg	acc	cat	ctc	atg	ggt	ctc	ttt	gag	cct	gga	gac	atg	aaa	tac	gag	6439
	Val	Thr	His	Leu	Met	Gly	Leu	Phe	Glu	Pro	Gly	Asp	Met	Lys	Tyr	Glu	
		280					285					290					
70	atc	cac	cga	gac	tcc	aca	ctg	gac	ccc	tcc	ctg	atg	gag	atg	aca	gag	6487
	Ile	His	Arg	Asp	Ser	Thr	Leu	Asp	Pro	Ser	Leu	Met	Glu	Met	Thr	Glu	
		295				300					305					310	
75	gct	gcc	ctg	cgc	ctg	ctg	agc	agg	aac	ccc	cgc	ggc	ttc	ttc	ctc	ttc	6535
	Ala	Ala	Leu	Arg	Leu	Leu	Ser	Arg	Asn	Pro	Arg	Gly	Phe	Phe	Leu	Phe	
					315					320					325		

	gtg gag ggt ggt cgc atc gac cat ggt cat cat gaa agc agg gct tac	6583
	Val Glu Gly Gly Arg Ile Asp His Gly His His Glu Ser Arg Ala Tyr	
	330 335 340	
5	cgg gca ctg act gag acg atc atg ttc gac gac gcc att gag agg gcg	6631
	Arg Ala Leu Thr Glu Thr Ile Met Phe Asp Asp Ala Ile Glu Arg Ala	
	345 350 355	
10	ggc cag ctc acc agc gag gag gac acg ctg agc ctc gtc act gcc gac	6679
	Gly Gln Leu Thr Ser Glu Glu Asp Thr Leu Ser Leu Val Thr Ala Asp	
	360 365 370	
15	cac tcc cac gtc ttc tcc ttc gga ggc tac ccc ctg cga ggg agc tcc	6727
	His Ser His Val Phe Ser Phe Gly Gly Tyr Pro Leu Arg Gly Ser Ser	
	375 380 385 390	
	atc ttc ggg ctg gcc cct ggc aag gcc cgg gac agg aag gcc tac acg	6775
	Ile Phe Gly Leu Ala Pro Gly Lys Ala Arg Asp Arg Lys Ala Tyr Thr	
	395 400 405	
20	gtc ctc cta tac gga aac ggt cca ggc tat gtg ctc aag gac ggc gcc	6823
	Val Leu Leu Tyr Gly Asn Gly Pro Gly Tyr Val Leu Lys Asp Gly Ala	
	410 415 420	
25	cgg ccg gat gtt acc gag agc gag agc ggg agc ccc gag tat cgg cag	6871
	Arg Pro Asp Val Thr Glu Ser Glu Ser Gly Ser Pro Glu Tyr Arg Gln	
	425 430 435	
30	cag tca gca gtg ccc ctg gac gaa gag acc cac gca ggc gag gac gtg	6919
	Gln Ser Ala Val Pro Leu Asp Glu Glu Thr His Ala Gly Glu Asp Val	
	440 445 450	
35	gcg gtg ttc gcg cgc ggc ccg cag gcg cac ctg gtt cac ggc gtg cag	6967
	Ala Val Phe Ala Arg Gly Pro Gln Ala His Leu Val His Gly Val Gln	
	455 460 465 470	
	gag cag acc ttc ata gcg cac gtc atg gcc ttc gcc gcc tgc ctg gag	7015
	Glu Gln Thr Phe Ile Ala His Val Met Ala Phe Ala Ala Cys Leu Glu	
	475 480 485	
40	ccc tac acc gcc tgc gac ctg gcg ccc ccc gcc ggc acc acc gac gcc	7063
	Pro Tyr Thr Ala Cys Asp Leu Ala Pro Pro Ala Gly Thr Thr Asp Ala	
	490 495 500	
45	gcg cac ccg ggt tac tct aga gtc ggg gcg gcc ggc cgc ttc gag cag	7111
	Ala His Pro Gly Tyr Ser Arg Val Gly Ala Ala Gly Arg Phe Glu Gln	
	505 510 515	
50	aca tga taagatacat tgatgagttt ggacgcggcc gcgactctag atcataatca	7167
	Thr	
	gccataccac atttgtagag gttttacttg ctttaaaaaa cctccacac ctccccctga	7227
55	acctgaaaca taaaatgaat gcaattgttg ttgttaactt gtttattgca gottataatg	7287
	gttacaaata aagcaatagc atcaciaaatt tcacaaataa agcatttttt tcaactgcatt	7347
	ctagttgtgg tttgtccaaa ctcatcaatg tatcttaagg cgtaaattgt aagcggttaat	7407
60	attttggttaa aattcgcggt aaatttttgt taaatcagct catttttttaa ccaataggcc	7467
	gaaatcgcca aaatccctta taaatcaaaa gaatagaccg agatagggtt gagtgttggt	7527

	ccagtttggga acaagagtcc actatttaaag aacgtggact ccaacgtcaa agggcgaaaa	7587
	accgtctatc agggcgatgg cccactacgt gaaccatcac cctaatacaag ttttttgggg	7647
5	tcgaggtgcc gtaaagcact aaatcggaac cctaaaggga gccccgatt tagagcttga	7707
	cggggaaagc cggcgaaact ggcgagaaag gaagggaaga aagcgaaagg agcgggagct	7767
10	agggcgctgg caagtgtagc ggtcacgctg cgcgtaacca ccacacccgc cgcgcttaat	7827
	gcgcgcgtac agggcgcgctc aggtggcact tttcggggaa atgtgcgcgg aaccctatt	7887
15	tgtttatattt tctaaataca ttcaaataatg tatccgctca tgagacaata accctgataa	7947
	atgcttcaat aatattgaaa aaggaagagt cctgaggcgg aaagaaccag ctgtggaatg	8007
	tgtgtcagtt aggggtgtgga aagtccccag gctccccagc aggcagaagt atgcaaagca	8067
20	tgcattctcaa ttagtcagca accaggtgtg gaaagtcccc aggctcccca gcaggcagaa	8127
	gtatgcaaag catgcatctc aattagtcag caaccatagt cccgccccta actccgcca	8187
	tcccgcctt aactccgcc agttccgcc atttccgcc ccatggctga ctaatttttt	8247
25	ttatttatgc agaggccgag gccgcctcgg cctctgagct attccagaag tagtgaggag	8307
	gcttttttgg aggcctaggc ttttgcaaag atcgatcaag agacaggatg aggatcgttt	8367
30	cgcattgattg aacaagatgg attgcacgca ggttctccgg ccgcttgggt ggagaggcta	8427
	ttcggctatg actgggcaca acagacaatc ggctgctctg atgccgccgt gttccggctg	8487
	tcagcgcagg ggcgcccggg tctttttgtc aagaccgacc tgtccggtgc cctgaatgaa	8547
35	ctgcaagacg aggcagcgcg gctatcgtgg ctggccacga cgggcgttcc ttgcgcagct	8607
	gtgctcgacg ttgtcactga agcgggaagg gactggctgc tattgggcca agtgccgggg	8667
40	caggatctcc tgtcatctca ccttgctcct gccgagaaag tatccatcat ggctgatgca	8727
	atgcggcggc tgcatacgtc tgatccggct acctgcccac tcgaccacca agcgaaacat	8787
	cgcattcgagc gagcacgtac tcggatggaa gccggtcttg tcgatcagga tgatctggac	8847
45	gaagagcatc aggggctcgc gccagccgaa ctgttcgcca ggctcaaggc gagcatgccc	8907
	gacggcgagg atctcgtcgt gacccatggc gatgcctgct tgccgaatat catggtggaa	8967
50	aatggccgct tttctggatt catcgactgt ggccggctgg gtgtggcgga ccgctatcag	9027
	gacatagcgt tggctacccg tgatattgct gaagagcttg gcggcgaatg ggctgaccgc	9087
	ttcctcgtgc ttacgggtat cgccgctccc gattcgcagc gcattgcctt ctatcgctt	9147
55	cttgacgagt tcttctgagc gggactctgg ggttcgaaat gaccgaccaa gcgacgcca	9207
	acctgccatc acgagatttc gattccaccg ccgccttcta tgaaagggtg ggcttcggaa	9267
60	tcgttttccg ggacgccggc tggatgatcc tccagcgcgg ggaatctcatg ctggagttct	9327
	tcgcccaccc tagggggagg ctaactgaaa cacggaagga gacaataacc gaaggaaccc	9387

gcgctatgac ggcaataaaa agacagaata aaacgcacgg tgttgggtcg tttgttcata 9447
 aacgcgggggt tcgggtcccag ggctggcact ctgtcgatac cccaccgaga cccattggg 9507
 5 gccaatacgc ccgcgtttct tctttttccc cccccaccc cccaagttcg ggtgaaggcc 9567
 cagggctcgc agccaacgtc ggggcggcag gccctgccat agcctcaggt tactcatata 9627
 10 tacttttagat tgatttaaaa cttcattttt aattttaaag gatctaggtg aagatccttt 9687
 ttgataatct catgaccaa atcccttaac gtgagttttc gttccactga gcgtcagacc 9747
 ccgtagaaaa gatcaaagga tcttcttgag atcctttttt tctgcgcgta atctgctgct 9807
 15 tgcaaacaaa aaaaccaccg ctaccagcgg tggtttgttt gccggatcaa gagctaccaa 9867
 ctctttttcc gaaggtaact ggcttcagca gagcgcagat accaaatact gtccttctag 9927
 tgtagccgta gttaggccac cacttcaaga actctgtagc accgcctaca tacctcgctc 9987
 20 tgctaatacct gttaccagtg gctgctgcca gtggcgataa gtcgtgtctt accgggttg 10047
 actcaagacg atagttaccg gataaggcgc agcggtcggg ctgaacgggg ggttcgtgca 10107
 25 cacagcccag cttggagcga acgacctaca ccgaactgag atacctacag cgtgagctat 10167
 gagaaagcgc cacgcttccc gaaggagaa aggcggacag gtatccggtg agcggcaggg 10227
 tcggaacagg agagcgcacg agggagcttc cagggggaaa cgcttggtat ctttatagtc 10287
 30 ctgtcggggt tcgccacctc tgacttgagc gtcgattttt gtgatgctcg tcaggggggc 10347
 ggagcctatg gaaaaacgcc agcaacgcgg cttttttacg gttcctggcc ttttgctggc 10407
 35 cttttgctca catgttcttt cctgcgttat cccctgattc tgtggataac cgtattaccg 10467
 ccatgcat 10475

 40 <210> 4
 <211> 519
 <212> PRT
 45 <213> SEAP

 50 <400> 4
 Met Leu Leu Leu Leu Leu Leu Gly Leu Arg Leu Gln Leu Ser Leu
 1 5 10 15
 55 Gly Ile Ile Pro Val Glu Glu Glu Asn Pro Asp Phe Trp Asn Arg Glu
 20 25 30
 60 Ala Ala Glu Ala Leu Gly Ala Ala Lys Lys Leu Gln Pro Ala Gln Thr
 35 40 45

Ala Ala Lys Asn Leu Ile Ile Phe Leu Gly Asp Gly Met Gly Val Ser
50 55 60

5 Thr Val Thr Ala Ala Arg Ile Leu Lys Gly Gln Lys Lys Asp Lys Leu
65 70 75 80

10 Gly Pro Glu Ile Pro Leu Ala Met Asp Arg Phe Pro Tyr Val Ala Leu
85 90 95

15 Ser Lys Thr Tyr Asn Val Asp Lys His Val Pro Asp Ser Gly Ala Thr
100 105 110

20 Ala Thr Ala Tyr Leu Cys Gly Val Lys Gly Asn Phe Gln Thr Ile Gly
115 120 125

25 Leu Ser Ala Ala Ala Arg Phe Asn Gln Cys Asn Thr Thr Arg Gly Asn
130 135 140

Glu Val Ile Ser Val Met Asn Arg Ala Lys Lys Ala Gly Lys Ser Val
145 150 155 160

30 Gly Val Val Thr Thr Thr Arg Val Gln His Ala Ser Pro Ala Gly Thr
165 170 175

35 Tyr Ala His Thr Val Asn Arg Asn Trp Tyr Ser Asp Ala Asp Val Pro
180 185 190

40 Ala Ser Ala Arg Gln Glu Gly Cys Gln Asp Ile Ala Thr Gln Leu Ile
195 200 205

Ser Asn Met Asp Ile Asp Val Ile Leu Gly Gly Gly Arg Lys Tyr Met
210 215 220

45 Phe Arg Met Gly Thr Pro Asp Pro Glu Tyr Pro Asp Asp Tyr Ser Gln
225 230 235 240

50 Gly Gly Thr Arg Leu Asp Gly Lys Asn Leu Val Gln Glu Trp Leu Ala
245 250 255

55 Lys Arg Gln Gly Ala Arg Tyr Val Trp Asn Arg Thr Glu Leu Met Gln
260 265 270

60 Ala Ser Leu Asp Pro Ser Val Thr His Leu Met Gly Leu Phe Glu Pro
275 280 285

Gly Asp Met Lys Tyr Glu Ile His Arg Asp Ser Thr Leu Asp Pro Ser
290 295 300

5 Leu Met Glu Met Thr Glu Ala Ala Leu Arg Leu Leu Ser Arg Asn Pro
 305 310 315 320

10 Arg Gly Phe Phe Leu Phe Val Glu Gly Gly Arg Ile Asp His Gly His
 325 330 335

15 His Glu Ser Arg Ala Tyr Arg Ala Leu Thr Glu Thr Ile Met Phe Asp
 340 345 350

20 Asp Ala Ile Glu Arg Ala Gly Gln Leu Thr Ser Glu Glu Asp Thr Leu
 355 360 365

25 Ser Leu Val Thr Ala Asp His Ser His Val Phe Ser Phe Gly Gly Tyr
 370 375 380

30 Pro Leu Arg Gly Ser Ser Ile Phe Gly Leu Ala Pro Gly Lys Ala Arg
 385 390 395 400

35 Asp Arg Lys Ala Tyr Thr Val Leu Leu Tyr Gly Asn Gly Pro Gly Tyr
 405 410 415

40 Val Leu Lys Asp Gly Ala Arg Pro Asp Val Thr Glu Ser Glu Ser Gly
 420 425 430

45 Ser Pro Glu Tyr Arg Gln Gln Ser Ala Val Pro Leu Asp Glu Glu Thr
 435 440 445

50 His Ala Gly Glu Asp Val Ala Val Phe Ala Arg Gly Pro Gln Ala His
 450 455 460

55 Leu Val His Gly Val Gln Glu Gln Thr Phe Ile Ala His Val Met Ala
 465 470 475 480

60 Phe Ala Ala Cys Leu Glu Pro Tyr Thr Ala Cys Asp Leu Ala Pro Pro
 485 490 495

65 Ala Gly Thr Thr Asp Ala Ala His Pro Gly Tyr Ser Arg Val Gly Ala
 500 505 510

70 Ala Gly Arg Phe Glu Gln Thr
 515

1. A method of monitoring cell differentiation comprising:
 - 5 (a) culturing cells capable of differentiating into at least one particular cell type containing at least one recombinant nucleic acid molecule comprising a reporter gene encoding a product that is secreted upon cell differentiation, or maintaining a non-human animal comprising such cells, under conditions allowing differentiation of the cells; and
 - 10 (b) determining the amount or activity of the reporter gene product either within a body fluid of said transgenic non-human animal or the cell culture medium.
- 15 2. The method of claim 1, wherein said recombinant nucleic acid molecule comprises at least one cell type specific regulatory sequence operably linked to said reporter gene.
3. The method of claim 1 or 2, wherein said cells are or are derived from stem cells or primordial germ cells, preferably human or rodent stem cells or primordial germ cells.
- 20 4. The method of any of claims 1 to 3, wherein said reporter gene product is secreted alkaline phosphatase (SEAP) or alpha-amylase.
- 25 5. The method of any one of claims 1 to 4, wherein said cells form embryoid bodies (EBs).
6. A reporter gene construct for monitoring cell differentiation comprising a recombinant nucleic acid molecule as defined in any one of claims 1 to 5.
- 30 7. A cell as defined in any one of claims 1 to 5 or comprising a reporter gene construct of claim 6, wherein said cell is capable of differentiating into at least one particular cell type.
8. A method of obtaining and/or profiling a modulator of cell differentiation comprising:

- (a) contacting a test sample comprising a cell of claim 7 with a test substance;
and
- (b) determining the effect of the test substance on the amount of the reporter gene product or activity compared to a control sample.

5

9. A kit useful for conducting a method of any one of claims 1 to 5 or 8, containing for example a reporter gene construct of claim 6 or a cell of claim 7, and standard compounds, like cell culture media, selection agents, detection agents for the reporter molecule and control samples.

10

10. A vector comprising the promoter region of the mouse alpha myosin heavy chain gene or of the ventricular myosin regulatory light chain gene, and operably linked thereto a heterologous DNA sequence, preferably wherein said promoter comprises the nucleotide sequence of SEQ ID NO: 1 or 2, or a fragment thereof.

15

ABSTRACT

EPO-Munich
08. Juli 2013

5 Provided are means and methods for in vitro and in vivo detection of chemically induced effects on embryonic development and differentiation for the purpose of embryotoxicity/teratogenicity screening as well as for the identification of pharmaceuticals such as growth and tissue promoting factors based on differentiating pluripotent embryonic stem (ES) cells. The assays are based on the use of transgenic ES cells and non-human animals comprising such ES cells or derivatives thereof, wherein said ES cells are characterized by the differentiation-dependent expression of a secreted reporter protein; in 10 particular secreted embryonic alkaline phosphatase (SEAP).




Fig. 1

